



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/57, 9/52, C07K 16/40, A01K 67/027, A61K 38/48, A61P 37/06		A1	(11) International Publication Number: WO 00/17370
			(43) International Publication Date: 30 March 2000 (30.03.00)
(21) International Application Number: PCT/US99/22362		Apartment 303, 319 8th Street Southeast, Minneapolis, MN 55414 (US). CHENG, Qi [CN/US]; 11240 48th Avenue North, Plymouth, MN 55442 (US). GREEN, Bruce, A. [US/US]; 49 Northfield Gate, Pittsford, NY 14534 (US). MASI, Amy, W. [US/US]; 326 Grand Circle, Caledonia, NY 14423 (US). (74) Agent: MUETING, Ann, M.; Mueeting, Raasch & Gebhardt, P.A., P.O. Box 581415, Minneapolis, MN 55458-1415 (US). (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 24 September 1999 (24.09.99)			
(30) Priority Data: 60/101,736 24 September 1998 (24.09.98) US 09/283,094 31 March 1999 (31.03.99) US			
(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Applications US 09/283,094 (CIP) Filed on 31 March 1999 (31.03.99) US 60/101,736 (CIP) Filed on 24 September 1998 (24.09.98)			
(71) Applicants (for all designated States except US): REGENTS OF THE UNIVERSITY OF MINNESOTA [US/US]; 100 Church Street SE, Minneapolis, MN 55455 (US). AMERICAN CYANAMID COMPANY [US/US]; 5 Giralda Farms, Madison, NJ 07940 (US).			
(72) Inventors; and (75) Inventors/Applicants (for US only): HOSTETTER, Margaret, K. [US/US]; Ezra Stiles College, 19 Tower Parkway, New Haven, CT 06511 (US). FINKEL, David, J. [US/US];		Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	
(54) Title: HUMAN COMPLEMENT C3-DEGRADING POLYPEPTIDE FROM STREPTOCOCCUS PNEUMONIAE			
<p>ATGTCAAGCCTTTTACGTGAATTGTATGCTAAACCCTTATCAGAACGCCATGTAGAATCTGATGCC TTATTTTCGACCCAGCGCAAATCACAAGTCGAACCGCCAATGGTGTTGCTGTACCGCACGGAGACC ATTATCACTTTATTCTTTATTCACAACTGTCACCTTTGGAAGAAAAATTGGCTCGTATTATTTCCCTTC GTTATCGTTCAAACCATTTGGGTACCAAGATTCAAGACCAGAACAACCAAGTCCACAATCGACTCCGG AACCTAGTCCAAGTCCGCAACCTGCACCAATCCTCAACCAGCTCCAAGCAATCCAATTGATGAGA AATTGGTCAAAGAAGCTGTTGAAAAAGTAGGCGATGGTTATGTCTTTGAGGAGAATGGAGTTCCTC GTTATATCCCAGCCAAGGATCTTTCAGCAGAAACAGCAGCAGGCATTGATAGCAAACCTGGCCAAGC AGGAAAGTTTATCTCATAAGCTGCAGTTAGATCCATTA</p>			
(57) Abstract			
<p>The present invention relates to the identification and use of a family of human complement C3-degrading polypeptides expressed by <i>S.pneumoniae</i>. The polypeptides have molecular weights of about 15 kD to about 25 kD or about 75 kDa to about 95 kDa, for example. Preferred polypeptides of this invention include the amino acid sequences of SEQ ID NO:2 and SEQ ID NO:5.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

HUMAN COMPLEMENT C3-DEGRADING POLYPEPTIDE FROM *STREPTOCOCCUS PNEUMONIAE*

5

Field of the Invention

This invention relates to *Streptococcus pneumoniae* and in particular this invention relates to the identification of an *S. pneumoniae* polypeptide that is capable of degrading human complement protein C3.

10

Background of the Invention

Respiratory infection with the bacterium *Streptococcus pneumoniae* (*S. pneumoniae*) leads to an estimated 500,000 cases of pneumonia and 47,000 deaths annually. Those persons at highest risk of bacteremic pneumococcal infection are infants under two years of age, individuals with a compromised immune system and the elderly. In these populations, *S. pneumoniae* is the leading cause of bacterial pneumonia and meningitis. Moreover, *S. pneumoniae* is the major bacterial cause of ear infections in children of all ages. Both children and the elderly share defects in the synthesis of protective antibodies to pneumococcal capsular polysaccharide after either bacterial colonization, local or systemic infection, or vaccination with purified polysaccharides. *S. pneumoniae* is the leading cause of invasive bacterial respiratory disease in both adults and children with HIV infection and produces hematogenous infection in these patients (Connor et al. *Current Topics in AIDS* 1987;1:185-209 and Janoff et al. *Ann. Intern. Med.* 1992;117(4):314-324).

Individuals who demonstrate the greatest risk for severe infection are not able to make antibodies to the current capsular polysaccharide vaccines. As a result, there are now four conjugate vaccines in clinical trial. Conjugate vaccines consist of pneumococcal capsular polysaccharides coupled to protein carriers or adjuvants in an attempt to boost the antibody response. However, there are other potential problems with conjugate vaccines currently in clinical trials. For example, pneumococcal serotypes that are most prevalent in the United States are different from the serotypes that are most common in places

30

such as Israel, Western Europe, South Africa, or Scandinavia. Therefore, vaccines that may be useful in one geographic locale may not be useful in another. The potential need to modify currently available capsular polysaccharide vaccines or to develop protein conjugates for capsular vaccines to suit geographic serotype variability entails prohibitive financial and technical complications. Thus, the search for immunogenic, surface-exposed proteins that are conserved worldwide among a variety of virulent serotypes is of prime importance to the prevention of pneumococcal infection and to the formulation of broadly protective pneumococcal vaccines. Moreover, the emergence of penicillin and cephalosporin-resistant pneumococci on a worldwide basis makes the need for effective vaccines even more exigent (Baquero et al. *J. Antimicrob. Chemother.* 1991;28S:31-8).

Several pneumococcal proteins have been proposed for conjugation to pneumococcal capsular polysaccharide or as single immunogens to stimulate immunity against *S. pneumoniae*. Surface proteins that are reported to be involved in adhesion of *S. pneumoniae* to epithelial cells of the respiratory tract include PsaA, PspC/CBP112, and IgA1 proteinase (Sampson et al. *Infect. Immun.* 1994;62:319-324, Sheffield et al. *Microb. Pathogen.* 1992; 13: 261-9, and Wani, et al. *Infect. Immun.* 1996; 64:3967-3974). Antibodies to these adhesins could inhibit binding of pneumococci to respiratory epithelial cells and thereby reduce colonization. Other cytosolic pneumococcal proteins such as pneumolysin, autolysin, neuraminidase, or hyaluronidase are proposed as vaccine antigens because antibodies could potentially block the toxic effects of these proteins in patients infected with *S. pneumoniae*. However, these proteins are typically not located on the surface of *S. pneumoniae*, rather they are secreted or released from the bacterium as the cells lyse and die (Lee et al. *Vaccine* 1994; 12:875-8 and Berry et al. *Infect. Immun.* 1994; 62:1101-1108). While use of these cytosolic proteins as immunogens might ameliorate late consequences of *S. pneumoniae* infection, antibodies to these proteins would neither promote pneumococcal death nor prevent initial or subsequent pneumococcal colonization.

A prototypic surface protein that is being tested as a pneumococcal

vaccine is the pneumococcal surface protein A (PspA). PspA is a heterogeneous protein of about 70-140 kDa. The PspA structure includes an alpha helix at the amino terminus, followed by a proline-rich sequence, and terminates in a series of 11 choline-binding repeats at the carboxy-terminus. Although much
5 information regarding its structure is available, PspA is not structurally conserved among a variety of pneumococcal serotypes, and its function is entirely unknown (Yother et al. *J. Bacteriol.* 1992;174:601-9 and Yother *J. Bacteriol.* 1994;176:2976-2985). Studies have confirmed the immunogenicity of PspA in animals (McDaniel et al. *Microb. Pathogen.* 1994; 17:323-337).
10 Despite the immunogenicity of PspA, the heterogeneity of PspA, its existence in four structural groups (or clades), and its uncharacterized function complicate its ability to be used as a vaccine antigen.

In patients who cannot make protective antibodies to the type-specific polysaccharide capsule, the third component of complement, C3, and the
15 associated proteins of the alternative complement pathway constitute the first line of host defense against *S. pneumoniae* infection. Because complement proteins cannot penetrate the rigid cell wall of *S. pneumoniae*, deposition of opsonic C3b on the pneumococcal surface is the principal mediator of pneumococcal clearance. Interactions of pneumococci with plasma C3 are
20 known to occur during pneumococcal bacteremia, when the covalent binding of C3b, the opsonically active fragment of C3, initiates phagocytic recognition and ingestion (Johnston et al. *J. Exp. Med.* 1969;129:1275-1290, Hasin HE, *J. Immunol.* 1972; 109:26-31 and Hostetter et al. *J. Infect. Dis.* 1984; 150:653-61). C3b deposits on the pneumococcal capsule, as well as on the cell wall. This
25 method for controlling *S. pneumoniae* infection is fairly inefficient. Methods for augmenting *S. pneumoniae* opsonization could improve the disease course induced by this organism. There currently exists a strong need for methods and therapies to limit *S. pneumoniae* infection.

30

Summary of the Invention

This invention relates to the identification and use of a family of human

complement C3-degrading polypeptides (e.g., proteins) expressed by *S. pneumoniae*. The polypeptides preferably have a molecular weight of about 15 kDa to about 25 kDa, or about 75 kDa to about 95 kDa, as determined, for example, on a 10% SDS polyacrylamide gel. The invention includes a number
5 of polypeptides isolatable from different C3-degrading strains of *S. pneumoniae*.

In one aspect, the invention relates to an isolated polypeptide having at least 80% sequence identity with SEQ ID NO:2 or SEQ ID NO:5. In a preferred embodiment, the polypeptide is isolated from *S. pneumoniae* or alternatively the polypeptide is a recombinant polypeptide. Preferably, the isolated polypeptide
10 degrades human complement protein C3. A preferred polypeptide of this invention is an isolated polypeptide having an amino acid sequence that includes SEQ ID NO:2 or SEQ ID NO:5, and more preferably, is SEQ ID NO:2 or SEQ ID NO:5. The term "isolated" as used herein refers to a naturally occurring species that has been removed from its natural environment, as well as to
15 synthetic species. The term "polypeptide" as used herein includes peptides, polypeptides, and proteins, regardless of length. Preferably, the polypeptides of the invention include one or more functional units, which encompasses polypeptides that degrade human complement protein C3.

Thus, the invention also relates to polypeptide fragments isolated from a
20 C3-degrading protein of this invention. Such fragments are encompassed by the term "polypeptide" as used herein. Preferably, the invention provides polypeptides of at least 15 sequential amino acids derived from a protein that has at least 80% sequence identity with SEQ ID NO:2 or SEQ ID NO:5, and more preferably, polypeptides of at least 15 sequential amino acids of SEQ ID NO:2
25 or SEQ ID NO:5. In another aspect of this invention, preferred polypeptides are capable of degrading human complement protein C3.

In another aspect, the invention relates to an isolated polypeptide that preferably degrades human complement protein C3, wherein nucleic acid encoding the isolated polypeptide hybridizes to at least a portion of SEQ ID
30 NO:1 or SEQ ID NO:4 or their complementary strands under highly stringent hybridization conditions. Preferably, the polypeptide includes at least 15

sequential amino acids, which are, more preferably, of SEQ ID NO:2 or SEQ ID NO:5.

In an additional aspect, this invention relates to polypeptides that have reduced human C3 degradation activity or which do not degrade human C3; however, the nucleic acids encoding this group of polypeptides each include a nucleotide sequence that hybridizes to either the nucleic acids that encode human C3 degrading polypeptides, or the complementary strands for each nucleic acid. This latter group of polypeptides having reduced or no human C3 degrading activity are referred to herein as "non-degrading" polypeptides. The non-degrading polypeptides may differ from C3 degrading polypeptides by one or more amino acids. This amino acid change may be a substitution, alteration, or deletion of one or more amino acids. Various types of amino acid changes are discussed herein. Nucleic acids encoding the non-degrading polypeptides are alternatively preferred embodiments of this invention.

The invention also relates to an immune system stimulating composition (preferably, a vaccine) comprising an effective amount of an immune system stimulating polypeptide of the present invention, which is preferably isolated from *S. pneumoniae*, and a therapeutically acceptable carrier. In one embodiment, the immune system stimulating composition or vaccine further comprises at least one other immune system stimulating polypeptide isolated from *S. pneumoniae*.

The invention further relates to an antibody capable of binding (typically, specifically binding) to a polypeptide (at least a portion thereof) of the present invention. In one embodiment, the antibody is a monoclonal antibody and in another embodiment, the antibody is a polyclonal antibody. In another embodiment the antibody is an antibody fragment. The antibody or antibody fragments can be obtained from a mouse, a rat, a goat, a chicken, a human, or a rabbit.

The invention also relates to an isolated nucleic acid molecule (i.e., a polynucleotide, which can be single stranded or double stranded, and which can be a part, or fragment, of a larger molecule such as a vector) capable of hybridizing to at least a portion of SEQ ID NO:1 or SEQ ID NO:4 or their

complimentary strands under highly stringent hybridization conditions. As used herein, highly stringent hybridization conditions include, for example, 6XSSC, 5X Denhardt, 0.5% SDS, and 100 µg/ml fragmented and denatured salmon sperm DNA hybridized overnight at 65°C and washed in 2X SSC, 0.1% SDS one time at room temperature for about 10 minutes followed by one time at 65°C for about 15 minutes followed by at least one wash in 0.2XSSC, 0.1% SDS at room temperature for at least 3-5 minutes. In one embodiment, the nucleic acid molecule is isolated from *S. pneumoniae* and in another embodiment, the nucleic acid molecule encodes a polypeptide. In one embodiment, the polypeptide degrades human complement protein C3. In another embodiment, the nucleic acid molecule encodes a polypeptide that does not degrade human complement C3.

In another embodiment, the nucleic acid molecule is in a vector (i.e., it is a fragment of a nucleic acid vector). The vector can be an expression vector capable of producing a polypeptide. Cells containing the nucleic acid molecule are also contemplated in this invention. In one embodiment, the cell is a bacterium or a eukaryotic cell.

The invention further relates to an isolated nucleic acid molecule comprising the nucleic acid sequence of SEQ ID NO:1 or SEQ ID NO:4, or their complementary strands. The invention further relates to an RNA molecule transcribed by a double-stranded DNA sequence comprising SEQ ID NO:1 or SEQ ID NO:4.

In another aspect of this invention, the invention relates to a method for producing an immune response to *S. pneumoniae* in a mammal (particularly a human). The method includes: administering to a mammal a composition comprising a therapeutically effective amount of a polypeptide of the present invention, and a pharmaceutically acceptable carrier, to produce an immune response to the polypeptide. The immune response can be a B cell response, a T cell response, an epithelial response, or an endothelial response. In a preferred embodiment, the composition is a vaccine composition. Preferably the polypeptide is at least 15 amino acids in length and also preferably the composition further comprises at least one other immune system stimulating

polypeptide from *S. pneumoniae*. In one embodiment, the polypeptide comprises at least 15 amino acids of SEQ ID NO:2 or SEQ ID NO: 5.

The invention further relates to an isolated polypeptide of about 15 kDa to about 25 kDa, or about 75 kDa to about 95 kDa, from *Streptococcus pneumoniae* that is capable of degrading human complement C3 and to a method for inhibiting *Streptococcus pneumoniae*-mediated C3 degradation. The method includes contacting a *Streptococcus pneumoniae* bacterium with antibody capable of binding to a polypeptide (at least a portion thereof) of the present invention.

The invention also relates to a method for inhibiting C3-mediated inflammation and rejection in xenotransplantation. The method includes expressing on the surface of an organ of an animal used in xenotransplantation a polypeptide of the present invention. This method is particularly advantageous for causing, for example, the kidneys of pigs to express the polypeptide described herein and thereby to inhibit C3 mediated inflammation after xenotransplantation.

The invention also relates to an isolated nucleic acid molecule that contains a region of at least 15 nucleotides which hybridize under highly stringent hybridization conditions to at least a portion of a nucleic acid sequence of SEQ ID NO:1 or SEQ ID NO:4 or their complementary strands.

The invention also relates to isolated DNA molecules or primers having the nucleic acid sequences as shown in SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8 AND SEQ ID NO:9.

25

Brief Description of the Figures

Figure 1 provides the nucleic acid sequence of the translated portion of a C3-degrading polypeptide (approximately 20 kDa) gene of this invention (SEQ ID NO:1).

Figure 2 provides the amino acid sequence of a C3-degrading polypeptide (approximately 20 kDa) of this invention (SEQ ID NO:2).

Figure 3 diagrams the amino acid sequence of a C3-degrading polypeptide (approximately 20 kDa) - sitioned with the nucleic acid sequence

(double stranded) encoding a C3-degrading polypeptide according to this invention (SEQ ID NOS:1-3 wherein SEQ ID NO:3 is the complement of SEQ ID NO:1).

Figure 4 provides the nucleic acid sequence for a predicted 92 kDa amino acid sequence (SEQ ID NO:4).

Figure 5 provides the predicted 92 kDa amino acid sequence (SEQ ID NO:5).

Figure 6 shows sequence alignments of SEQ ID NO:1 and a portion of SEQ ID NO:4.

Figure 7 shows sequence alignment of SEQ ID NO:2 with a portion of SEQ ID NO:5.

Figure 8. Western blot analysis of several pneumococcal whole cell lysate with polyclonal anti-r20 kDa sera. Molecular weight markers were run in lane 1; recombinant 20 kDa polypeptide from pDF122 was run in lane 2; recombinant 92 kDa polypeptide from Type 7 was run in lane 3; lane 4 was a blank; CP1200 whole cell lysate was run in lane 5; Type 3 whole cell lysate was run in lane 6; and Type 7 whole cell lysate was run in lane 7. The antisera recognizes the recombinant polypeptides of approximately 20 kDa and approximately 92 kDa, but only recognized the larger polypeptide in whole cell lysates.

Figure 9. Autoradiogram showing degradation of biotinylated C3 by 20 kDa and 92 kDa polypeptides of the present invention.

Figure 10. Coomassie Blue-Stained 7.5% SDS-PAGE analysis of C3 degradation by the 20 kDa (Lane A8) and the 92 kDa (Lane C8) polypeptides of the present invention.

Figure 11 is a chart which shows the survival of CBA/CAHN *xid*^{-/-} mice immunized SC (subcutaneous) with r79 kDa protein adjuvanted with MPL and challenged IN (intranasal) with *S. pneumoniae* Type 3.

Figure 12 is a SDS-PAGE gel described in Example 8.

Detailed Description of the Preferred Embodiments

The present invention relates to the identification and isolation of a human complement C3 degrading polypeptide fragment of a larger polypeptide of about 75 kD to about 95 kD. This fragment has a molecular weight of about 20 kDa (± 5 kDa) on a 10% SDS-PAGE gel. It also relates to nucleic acids encoding C3 degrading polypeptides.

It has been observed that exponentially growing cultures of pneumococci from several serotypes were able to first degrade the β -chain then degrade the α chain of C3 without producing defined C3 cleavage fragments (Angel, et al. *J. Infect. Dis.* 170:600-608, 1994). This pattern of degradation without cleavage differs substantially from that of other microbial products such as the elastase enzyme of *Pseudomonas aeruginosa* and the cysteine proteinase of *Entamoeba histolytica*.

The term "degrade" is used herein to refer to the removal of amino acids from proteinaceous molecules, generating peptides or polypeptides. The polypeptide of this invention degrades C3 without generating the cleavage fragments known as C3b, iC3b, or C3d. There is at least some preference of the C3-degrading polypeptides of this invention for C3 in that, for example, the C3-degrading polypeptide does not appear to degrade other proteins, such as albumin.

A C3-degrading polypeptide of about 20 kDa was isolated from a library of insertionally interrupted pneumococcal genes by identifying those clones that had decreased C3 degrading activity as compared to wild type *S. pneumoniae*. An exemplary assay for assessing C3-degrading activity of clones is provided in Example 1. Clones with decreased C3-degrading activity were identified and a 546 bp SmaI insert was selected, based on the sequence of the clones that had demonstrated decreased C3-degrading activity. This SmaI fragment was used to probe an *S. pneumoniae* library made from strain CP1200. Positive clones from the *S. pneumoniae* library that hybridized to the SmaI fragment were isolated and the open reading frame of the gene associated with C3-degrading activity was identified. The following oligonucleotide (SEQ ID NO:10), which has sequence identify with a portion of PspA, was used to confirm, by differential

hybridization, that the gene encoding the C3-degrading protinease was distinct from the gene encoding PspA.

SEQ ID NO:10

5 GAAAACAATAATGTAGAAGACTACTTTAAAGAAGGTTAGA

An open reading frame of a 20 kDa polypeptide spans an area of approximately 500 base pairs (SEQ ID NO:1) predicting a polypeptide of molecular weight of about 20 kDa (+/- 5 kDa) or about 168 amino acids (SEQ
10 ID NO:2). An exemplary gene sequence encoding a C3-degrading polypeptide is provided in Figure 1 as SEQ ID NO:1 and an amino acid sequence of the polypeptide is provided in Figure 2 as SEQ ID NO:2. Figure 3 combines a preferred gene sequence with a corresponding preferred translated polypeptide as SEQ ID NOS:1-3.

15 Using SEQ ID NO:2, the amino acid sequence of the approximately 20 kDa polypeptide, was determined to be unrelated to other polypeptides in the GenBank or Swiss Prot databases. The predicted polypeptide encompasses a proline-rich sequence characteristic of membrane domains in prokaryotes, particularly between amino acids 80-108 suggesting that the polypeptide is
20 expressed at the surface. The amino acid sequence exhibits no apparent choline-binding repeats. Electrophoresis of pneumococcal lysates and supernatants from cultures of CP 1200 on SDS-PAGE gels impregnated with C3 identified a lytic band at about 20 kDa (\pm 5 kDa) in both supernatants and lysates, confirming that a polypeptide of a size predicted by SEQ ID NO:2 had C3-degrading activity
25 (see Example 2). As provided in Example 3, the gene encoding the 20 kDa C3-degrading polypeptide is conserved in at least two dozen pneumococcal isolates representing five serotypes (serotypes 1, 3, 4, 14, and 19F).

The nucleotide sequence encoding the C3-degrading polypeptide of this invention was inserted into a gene expression vector for expression in *E. coli*.
30 Recombinant C3-degrading polypeptide was isolated as described in the examples. Those of ordinary skill in the art recognize that, given a particular gene sequence such as that provided in SEQ ID NO:1, there are a variety of

expression vectors that could be used to express the gene. Further, there are a variety of methods known in the art that could be used to produce and isolate the recombinant polypeptide of this invention and those of ordinary skill in the art also recognize that the C3 degrading assay of this invention will determine
5 whether or not a particular expression system, in addition to those expression systems provided in the examples, is functioning, without requiring undue experimentation. A variety of molecular and immunological techniques can be found in basic technical texts such as those of Sambrook et al. (*Molecular Cloning, A Laboratory Manual*, 1989 Cold Spring Harbor Laboratory Press,
10 Cold Spring Harbor, NY) and Harlow et al. (*Antibodies; A Laboratory Manual*. Cold Spring Harbor, NY; Cold Spring Harbor Laboratory Press, 1988).

The polynucleotide encoding the C3 degrading polypeptide of this invention was identified using a plasmid library made with pneumococcal genomic DNA fragments from strain CP1200. Although there are a variety of
15 methods known for obtaining a plasmid library, in a preferred strategy, a plasmid library was constructed with Sau 3A digested pneumococcal genomic DNA fragments (0.5 -4.0 kb) from pneumococcal strain CP 1200 (obtained from D.A. Morrison, University of Illinois, Champagne-Urbana, Illinois and described in Havarstein LF, et al. *Proc. Natl. Acad. Sci. (USA)* 1995;92:11140-11144) and
20 inserted into the Bam HI site of the integrative shuttle vector pVA 891 (*erm*^r, *cm*^r; has origin of replication for *E. coli*). This library was transformed into an *E. coli* DH5 α MCR strain by electroporation. Plasmid extractions of some randomly selected *E. coli* transformants revealed that all of them contained recombinant plasmids.

25 Plasmid library DNA can be extracted from the *E. coli* transformants and used to transform the CP 1200 parent pneumococcal strain using insertional mutagenesis by homologous recombination.

The pneumococcal strain CP 1200 cells can be made competent using a pH shift with HCl procedure in CTM medium. The competent cells are frozen at
30 -70°C in small aliquots until needed.

The isolated polypeptide of this invention can be incubated with human complement C3 for 4 hours or longer at 37°C in the presence of PBS to detect

C3 degradation. Control samples without the isolated pneumococcal polypeptide are used as controls for comparative purposes.

The polypeptides of this invention have an apparent molecular weight on a 10% SDS-polyacrylamide gel of either about 15 kDa to about 25 kDa; and preferably about 20 kDa; or about 75 kDa to about 95 kDa; and preferably about 92 kDa. Exemplary polypeptides sequences are provided by SEQ ID NO: 2 and SEQ ID NO:5. Those of ordinary skill in the art will recognize that some variability in amino acid sequence is expected and that this variability should not detract from the scope of this invention. For example, conserved mutations do not detract from this invention nor do variations in amino acid sequence identity of less than about 80% amino acid sequence identity and preferably less than about 90% amino acid sequence identity where the polypeptide is capable of degrading human complement protein C3, and particularly where the polypeptide is isolated or originally obtained from an *S. pneumoniae* bacterium. Proteins and fragments thereof (all referred to as polypeptides) are also within the scope of the present invention, particularly if they are capable of degrading human complement protein C3.

Some nucleic acid sequence variability is expected among pneumococcal strains and serotypes as is some amino acid variability. Conserved amino acid substitutions are known in the art and include, for example, amino acid substitutions using other members from the same class to which the amino acid belongs. For example, the nonpolar amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, and tryptophan. The polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine and glutamine. The positively charged (basic) amino acids include arginine, lysine and histidine. The negatively charged (acidic) amino acids include aspartic acid and glutamic acid. Such alterations are not expected to affect apparent molecular weight as determined by polyacrylamide gel electrophoresis or isoelectric point. Particularly preferred conservative substitutions include, but are not limited to, Lys for Arg and *vice versa* to maintain a positive charge; Glu for Asp and *vice versa* to maintain a negative charge; Ser for Thr so that a free -OH is maintained; and Gln for Asn to maintain a free NH₂.

Preferred polypeptides of this invention includes polypeptides with the amino acid sequence of SEQ ID NO:2 or SEQ ID NO:5. Other polypeptides include those degrading human complement polypeptide C3 and having nucleic acid encoding the polypeptide that hybridizes to SEQ ID NO:1 or SEQ ID NO:4 under highly stringent hybridization conditions such as 6XSSC, 5X Denhardt, 0.5% SDS (sodium dodecyl sulfate), and 100 µg/ml fragmented and denatured salmon sperm DNA hybridized overnight at 65°C and washed in 2X SSC, 0.1% SDS one time at room temperature for about 10 minutes followed by one time at, 65°C for about 15 minutes followed by at least one wash in 0.2XSSC, 0.1% SDS at room temperature for at least 3-5 minutes are also contemplated in this invention. Typically, an SSC solution contains sodium chloride, sodium citrate, and water to prepare a stock solution.

The polypeptides of this invention can be isolated or prepared as recombinant polypeptides. That is, nucleic acid encoding a protein, or a portion thereof, can be incorporated into an expression vector or incorporated into a chromosome of a cell to express the polypeptide in the cell. The polypeptide can be purified from a bacterium or another cell, preferably a eukaryotic cell and more preferably an animal cell. Alternatively, the polypeptide can be isolated from a cell expressing the polypeptide, such as a *S. pneumoniae* cell. Thus, proteins, peptides, or polypeptides are all considered within the scope of this invention when the term "polypeptide" is used. The polypeptides are preferably at least 15 amino acids in length and preferred polypeptides have at least 15 sequential amino acids from SEQ ID NO:2 or SEQ ID NO:5.

Nucleic acid encoding the about 15 kDa to about 20 kDa polypeptide and the about 75 kD to about 95 kD polypeptide are also part of this invention. SEQ ID NOS:1 and 4 are preferred nucleic acid molecules encoding a C3-degrading polypeptide. Those of ordinary skill in the art will recognize that some substitution will not alter the C3-degrading polypeptide sequence to an extent that the character or nature of the C3-degrading polypeptide is substantially altered. For example, nucleic acid with an identity of at least 80% to SEQ ID NO:1 is contemplated in this invention. A method for determining whether a particular nucleic acid sequence falls within the scope of this invention is to

consider whether or not a particular nucleic acid sequence encodes a C3-degrading polypeptide and has a nucleic acid identity of at least 80% as compared with SEQ ID NO:1 or SEQ ID NO:4. Other nucleic acid sequences encoding the C3 polypeptide include nucleic acid encoding the C3 polypeptide where the C3 polypeptide has the same sequence or at least a 90% sequence identity with SEQ ID NO:2 or SEQ ID NO:5 but which includes degeneracy with respect to the nucleic acid sequence. A degenerate codon means that a different three letter codon is used to specify the same amino acid. For example, it is well known in the art that the following RNA codons (and therefore, the corresponding DNA codons, with a T substituted for a U) can be used interchangeably to code for each specific amino acid:

	Phenylalanine (Phe or F)	UUU, UUC, UUA or UUG
	Leucine (Leu or L)	CUU, CUC, CUA or CUG
	Isoleucine (Ile or I)	AUU, AUC or AUA
15	Methionine (Met or M)	AUG
	Valine (Val or V)	GUU, GUC, GUA, GUG
	Serine (Ser or S)	AGU or AGC
	Proline (Pro or P)	CCU, CCC, CCA, CCG
	Threonine (Thr or T)	ACU, ACC, ACA, ACG
20	Alanine (Ala or A)	GCU, GCG, GCA, GCC
	Tryptophan (Trp)	UGG
	Tyrosine (Tyr or Y)	UAU or UAC
	Histidine (His or H)	CAU or CAC
	Glutamine (Gln or Q)	CAA or CAG
25	Asparagine (Asn or N)	AAU or AAC
	Lysine (Lys or K)	AAA or AAG
	Aspartic Acid (Asp or D)	GAU or GAC
	Glutamic Acid (Glu or E)	GAA or GAG
	Cysteine (Cys or C)	UGU or UGC
30	Arginine (Arg or R)	AGA or AGG
	Glycine (Gly or G)	GGU or GGC or GGA or GGG
	Termination codon	UAA, UAG or UGA

Further, a particular DNA sequence can be modified to employ the codons preferred for a particular cell type. For example, the preferred codon usage for *E. coli* is known, as are preferred codons for animals including humans. These changes are known to those of ordinary skill in the art and therefore these gene sequences are considered part of this invention. Other nucleic acid sequences include at least 15, and preferably, at least 30 nucleic acids in length from SEQ ID NO:1 or SEQ ID NO:4 or other nucleic acid fragments of at least 15, and preferably at least 30 nucleic acids in length where these fragments hybridize to SEQ ID NO:1 or SEQ ID NO:4 under highly stringent hybridization conditions such as 6XSSC, 5X Denhardt, 0.5% SDS, and 100 µg/ml fragmented and denatured salmon sperm DNA hybridized overnight at 65°C and washed in 2X SSC, 0.1% SDS one time at room temperature for about 10 minutes followed by one time at, 65°C for about 15 minutes followed by at least one wash in 0.2XSSC, 0.1% SDS at room temperature for at least 3-5 minutes.

The nucleic acid molecules of this invention can encode all, none (i.e., fragments that cannot be transcribed, fragments that include regulatory portions of the gene, or the like), or a portion of SEQ ID NO:2 or SEQ ID NO:5 and preferably containing a contiguous nucleic acid fragment that encodes at least nine amino acids from SEQ ID NO:2 or SEQ ID NO:5. Because nucleic acid molecules encoding a portion of a C3 degrading polypeptide are contemplated in this invention, it will be understood that not all of the nucleic acid molecules will encode a protein or peptide or polypeptide with C3 degrading activity. Further, the nucleic acid of this invention can be mutated to remove or otherwise inactivate the C3 degrading activity of this polypeptide. Therefore, nucleic acid molecules that encode polypeptides without C3 degrading activity that meet the hybridization requirements described above are also contemplated. Methods for mutating or otherwise altering nucleic acid sequences are well described in the art and the production of an immunogenic, but enzymatically inactive polypeptide can be tested for therapeutic utility.

The nucleic acid molecules of this invention can be incorporated into nucleic acid vectors or stably incorporated into host genomes to produce

recombinant polypeptides including recombinant chimeric polypeptides. In one embodiment, the C3-degrading polypeptide is encoded by a gene in a vector and the vector is in a cell. Preferably, the cell is a prokaryotic cell such as a bacterium. The genes and gene fragments can exist as the fusion of all or a portion of the gene with another gene and the C3-degrading polypeptide can exist as a fusion protein of one or more proteins where the fusion protein is expressed as a single protein. A variety of nucleic acid vectors of this invention are known in the art and include a number of commercially available expression plasmids or viral vectors. The use of these vectors is well within the scope of what is ordinary skill in the art. Exemplary vectors are employed in the examples, but should not be construed as limiting on the scope of this invention.

This invention also relates to antibodies capable of binding (typically specifically binding) to polypeptides of about 15 kDa to about 25 kDa; and preferably about 20 kDa; and about 75 kDa to about 95 kDa; and preferably about 92 kDa, from *S. pneumoniae* and preferably where the polypeptides are capable of degrading human complement C3. Polyclonal or monoclonal antibodies can be prepared to all or part of the polypeptides of the invention. Methods for preparing antibodies to polypeptides are well known and well described, for example, by Harlow et al., (*Antibodies; A Laboratory Manual*. Cold Spring Harbor, NY; Cold Spring Harbor Laboratory Press, 1988). In a preferred example, the antibodies can be human derived, rat derived, mouse derived, goat derived, chicken derived, or rabbit derived. Polypeptide-binding antibody fragments and chimeric fragments are also known and are within the scope of this invention.

The invention also relates to the use of immune stimulating compositions. The term "immune stimulating" or "immune system stimulating" composition refers to protein, peptide, or polypeptide compositions according to the invention that activate at least one cell type of the immune system in a subject, such as a mammal. Preferably, the immune stimulating composition provides an immunizing response or prophylactic benefit in a normal, i.e., uninfected subject, typically a vaccine. However, any measurable immune response is beneficial to the subject in a therapy application or protocol.

Preferred activated cells of the immune system include phagocytic cells such as neutrophils or macrophages, T cells, B cells, epithelial cells and endothelial cells. Immune stimulating compositions comprising the peptides, polypeptides or proteins of the invention can be used to produce antibody in an animal such as a rat, mouse, goat, chicken, rabbit, or a human or an animal model for studying *S. pneumoniae* infection. Preferred immune stimulating compositions include an immune stimulating amount, e.g., a therapeutically effective amount, of at least one peptide or polypeptide including at least 15 amino acids from the C3 degrading polypeptide.

10 The term "vaccine" refers to a composition for immunization. This process can include the administration of a protein, peptide, polypeptide, antigen, nucleic acid sequence or complementary sequence, e.g., anti-sense, or antibody, or suspensions thereof, wherein upon administration, the molecule will produce active immunity and provide protection against an *S. pneumoniae* infection or colonization. Typically, such vaccines are prepared as injectables, 15 either as liquid solutions or suspensions. Solid forms suitable for solution in, or suspension in, liquid prior to injection may also be prepared. The vaccine preparation may optionally be emulsified, or encapsulated in liposomes.

The immune stimulating composition (such as a vaccine) can further include other polypeptides in a pharmaceutically acceptable buffer or carrier, 20 such as PBS (phosphate buffer saline) or another buffer recognized in the art as suitable and safe for introduction of polypeptides into a host to stimulate the immune system. The immune stimulating compositions can also include other immune system stimulating polypeptides such as adjuvants or immune stimulating proteins, peptides, or polypeptides from *S. pneumoniae* or other 25 organisms. For example, a cocktail of peptides or polypeptides may be most useful for controlling *S. pneumoniae* infection. Preferably one or more of the polypeptides, or fragments thereof, of this invention are used in a vaccine preparation to protect against or limit *S. pneumoniae* colonization or the pathogenic consequences of *S. pneumoniae* colonization. 30

A "therapeutically effective amount," as used herein, refers to that amount that is effective for production of a desired result. This amount varies

depending upon the health and physical condition of a subject's immune system, i.e., to synthesize antibodies, the degree of protection desired, the formulation prepared and other relevant factors. It is expected that the amount will fall in a relatively broad range that can be determined through routine trials.

5 The active immune stimulating ingredients are often mixed with excipients or diluents that are pharmaceutically acceptable as carriers and compatible with the active ingredient. The term "pharmaceutically acceptable carrier" refers to a carrier(s) that is "acceptable" in the sense of being compatible with the other ingredients of a composition and not deleterious to the recipient
10 thereof. Suitable excipients are, for example, water, saline, dextrose, glycerol, ethanol, or the like and combinations thereof. In addition, if desired, the immune stimulating composition (including vaccine) may contain minor amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents, and/or adjuvants which enhance the effectiveness of the immune stimulating
15 composition.

Examples of adjuvants or carriers that may be effective include but are not limited to: aluminum hydroxide, N-acetyl-muramyl-L-threonyl-D-isoglutamine (thr-MDP); N-acetyl-nor-muramyl-L-alanyl-D-isoglutamine (CGP 11637, referred to as nor-MDP), N-acetylmuramyl-L-alanyl-D-isoglutaminyl-L-alanine-2-(1'-2'-dipalmitoyl-sn-glycero-3-hydroxyphosphoryloxy)-ethylamine
20 (CGP 19835A, referred to as MTP-PE), and RIBI, which contains three components extracted from bacteria, monophosphoryl lipid A, trehalose dimycolate and cell wall skeleton (MPL + TDM + CWS) in a 2% squalene/Tween 80 emulsion.

25 This invention also relates to a method for inhibiting *Streptococcus pneumoniae*-mediated C3 degradation comprising contacting a *Streptococcus pneumoniae* bacterium with a polypeptide, such as an antibody or another polypeptide that is capable of binding to an isolated polypeptide (typically, at least a portion thereof) of about 15 kDa to about 25 kDa, or about 75 kDa to
30 about 95 kDa, from *Streptococcus pneumoniae*. The protein capable of binding to an isolated polypeptide of about 15 kDa to about 25 kDa, or about 75 kDa to about 95 kDa can be an antibody or a fragment thereof, or the protein can be a

chimeric protein that includes the antibody binding domain, such as a variable domain, from antibody that is capable of specifically recognizing an isolated polypeptide of about 15 kDa to about 25 kDa, or about 75 kDa to about 95 kDa, from *Streptococcus pneumoniae* having C3 degrading activity.

5 The isolated *S. pneumoniae* polypeptide of this invention can be isolated, and optionally purified, and the isolated polypeptide or immunogenic fragments thereof can be used to produce an immunologic response, including, in one example, an antibody response in a human or an experimental animal. Polypeptides without C3 degrading ability can be tested for their ability to limit
10 the effects of *S. pneumoniae* infection. Similarly, the polypeptides of this invention can be modified, such as through mutation to interrupt or inactivate the C3 degrading activity of the polypeptides. Antibody capable of inhibiting the C3-degrading activity of the polypeptides of this invention may be used as a strategy for preventing C3 degradation and for promoting clearance of *S.*
15 *pneumoniae* through the opsonic pathway. Isolated polypeptides can be used in assays to detect antibody to *S. pneumoniae* or as part of a vaccine or a multi-valent or multiple protein, peptide, or polypeptide-containing vaccine for *S. pneumoniae* therapy.

 Thus, the term "treatment," as used herein, refers to prophylaxis and/or
20 therapy of either normal mammalian subjects or mammalian subjects colonized with, diagnosed with, or exhibiting characteristics or symptoms of various *S. pneumoniae* infections. The term "therapy" refers to providing a therapeutic effect to a mammalian subject such that the subject exhibits few or no symptoms of a pneumococcal infection or other related disease. Such treatment can be
25 accomplished by administration of nucleic acid molecules (sense or antisense), proteins, peptides or polypeptides or antibodies of the instant invention.

 It is further contemplated that the polypeptides of this invention can be surface expressed on vertebrate cells and used to degrade C3, for example, where complement deposition (or activation) becomes a problem, such as in
30 xenotransplantation or in complement-mediated glomerulonephritis. For example, the entire pneumococcal polypeptide, a recombinant polypeptide, or a portion of either, can be incorporated into xenotransplant cells and expressed as

a surface polypeptide or as a secreted polypeptide to prevent or minimize complement deposition (and/or complement-mediated inflammation).

Another specific aspect of the present invention relates to using a vaccine vector expressing an isolated protein, and peptides or polypeptides therefrom.

- 5 Accordingly, in a further aspect this invention provides a method of inducing an immune response in a mammal, which comprises providing to a mammal a vaccine vector expressing at least one, or a mixture of an isolated protein and/or peptide or polypeptide of the invention. The protein and peptides or polypeptides of the present invention can be delivered to the mammal using a
- 10 live vaccine vector, in particular using live recombinant bacteria, viruses or other live agents, containing the genetic material necessary for the expression of the protein and/or peptides or polypeptides as a foreign polypeptide. Particularly, bacteria that colonizes the gastrointestinal tract, such as *Salmonella*, *Shigella*, *Yersinia*, *Vibrio*, *Escherichia* and BCG have been developed as vaccine vectors,
- 15 and these and other examples are discussed by J. Holmgren et al., *Immunobiol.*, 184, 157-179 (1992) and J. McGhee et al., *Vaccine*, 10, 75-88 (1992).

- An additional embodiment of the present invention relates to a method of inducing an immune response in a subject, e.g., mammal, comprising administering to the subject an amount of a DNA molecule encoding an isolated
- 20 protein and/or peptide or polypeptide therefrom of this invention, optionally with a transfection-facilitating agent, where the protein and/or peptides or polypeptides retain immunogenicity and, when incorporated into an immune stimulating composition, e.g., vaccine, and administered to a human, provides protection without inducing enhanced disease upon subsequent infection of the
- 25 human with *S. pneumoniae* pathogen. Transfection-facilitating agents are known in the art.

- It is further contemplated that the antisense sequence of the gene encoding the about 15 kDa to about 25 kDa polypeptide, and the about 75 kDa to about 95 kDa polypeptide may be used as a vaccine or as a therapeutic treatment
- 30 for pneumococcal infection. Antisense DNA is defined as a non-coding sequence that is complementary, i.e., a complementary strand, to all or a portion of SEQ ID NO:1 or SEQ ID NO:4. For example, the antisense sequence for 5'-

ATGTCAAGC-3' is 3'-TACAGTTCG-5'. Delivery of antisense sequence or oligonucleotides into an animal may result in the production of antibody by the animal or in the incorporation of the sequence into living bacteria or other cells whereby transcription and/or translation of all or a portion of the 92 kDa gene product is inhibited.

Introduction of an antisense nucleic acid sequence can be accomplished, for example, by loading the antisense nucleic acid into a suitable carrier, such as a liposome, for introduction into pneumococci or infected cells. Typically, an antisense nucleic acid sequence having eight or more nucleotides is capable of binding to the bacterial nucleic acid or bacterial messenger RNA. The antisense nucleic acid sequence, typically contains at least about 15 nucleotides, preferably at least about 30 nucleotides or more nucleotides to provide necessary stability of a hybridization product of bacterial nucleic acid or bacterial messenger RNA. Introduction of the sequences preferably inhibit the transcription or translation of at least one endogenous *S. pneumoniae* nucleic acid sequence. Methods for loading antisense nucleic acid is known in the art as exemplified by U.S. Patent 4,242,046.

The present invention also provides nucleic acid having an open reading frame of 2478 bases (SEQ ID NO:4) that encompasses the open reading frame of a nucleic acid sequence (SEQ ID NO:1) that encodes a polypeptide that has a molecular weight of about 20 kDa (SEQ ID NO:2). The 20 kDa polypeptide, described herein, is further characterized as a C3-degrading polypeptide. The larger open reading frame, e.g., 2163 bp (SEQ ID NO:4), encodes for a putative polypeptide of about 92 kDa (SEQ ID NO:5).

All references and publications cited herein are expressly incorporated by reference into this disclosure. There are a variety of alternative techniques and procedures available to those of skill in the art which would similarly permit one to successfully perform the intended invention in view of the present disclosure. It will be appreciated by those skilled in the art that while the invention has been described above in connection with particular embodiments and examples, the invention is not necessarily so limited and that numerous other embodiments, examples, uses, modifications and departures from the embodiments, examples,

and uses may be made without departing from the inventive scope of this application.

5

Example 1

Identification of Insertional Mutants with Reduced C3-Degrading Activity

Insertional mutants were received from Dr. Elaine Tuomanen, (Rockefeller Inst., New York, New York). The clones with insertions were
10 tested in an assay to detect reduced C3-degrading activity. 137 clones were tested by growing the cells in Todd Hewitt broth overnight at room temperature in microtitre plates. The cells were diluted 1:10 in synthetic medium for pneumococci (see Sicard A. M., *Genetics* 50:31-44, 1984) and the remainder of the cells were frozen in the microtiter plate. Either 63 ng or 83 ng of C3
15 (purified from human plasma according to the method of Tack et al., *Meth. Enzymol.* 80:64-101, 1984) per 100 μ l of medium containing 1 mg/ml of 0.1% BSA in phosphate buffered saline (PBS) was added to about 200 μ l of diluted cells. The cells were incubated at 37°C for 4 hrs. One hundred μ l of the mixture was added to ELISA plates and incubated overnight at 4°C. The plates were
20 washed three times with wash buffer and the wells were filled with 0.05% Tween 20 in PBS with five minute incubations between the washes. One hundred μ l of antibody to C3 (polyclonal horse-radish peroxidase-conjugated goat antibody specific to human C3-IgG fraction, ICN Cappel, Costa Mesa, CA) was diluted 1:1200 with 3% BSA in PBS. The ELISA plate was incubated at
25 37°C for about 30 minutes to 1 hr in the dark and washed with wash buffer as above. The assay was developed using 12 mg of OPD in 30 ml of 0.1M sodium citrate buffer with 12 μ l of 30% hydrogen peroxide. Assay results were determined by optical density readings at 490 nm on an ELISA plate reader.

Each clone was tested four times. Nineteen clones were selected that had
30 less than 40% C3 degradation as compared to nonmutated controls. These 19 clones were screened 6 times by the assay described above and from these results 5 clones were selected with less than 30% C3-degrading activity as

compared to controls. These 6 clones were screened eleven times each and the two clones with the lowest C3-degrading activity were selected for further study.

A partial sequence of one of the clones was received and a *SmaI* fragment of 546 bp was labeled with ^{32}P by random primer labeling (kit available from Stratagene, La Jolla, CA). The 546bp *SmaI* fragment from SEQ ID NO:1 was hybridized to *EcoRI* and *KpnI* digests of numerous pneumococcal strains on Southern blots. This same fragment was also used to screen a library of *Sau3A* fragments of genomic DNA from *S. pneumoniae* strain CP1200.

A 3.5 kb insert was identified from the CP1200 library. The insert was sequenced and an open reading frame of 492 base pairs, including the stop codon, was identified. The open reading frame coded for a polypeptide of 168 amino acids and a predicted molecular weight of about 18,500 daltons.

PCR primers were constructed to amplify the open reading frame; the 5' PCR primer incorporated a *BamHI* site; the 3' primer incorporated a *PstI* site. The amplified insert was ligated in frame to a His-Tagged *E. coli* expression vector pQE30 (Qiagen, San Diego, CA). The resulting plasmid was used to transform *E. coli* strain BL21 (Novagen, Madison, WI) containing the lac repressor plasmid pREP4 (Qiagen). *E. coli* cultures were induced to express the His-Tagged polypeptide and the polypeptide was column purified with Ni-NTA resin (Qiagen). The purified polypeptide was confirmed by SDS-PAGE gel.

Example 2

Identification of a 20kDa C3-Degrading Polypeptide

To determine the C3-degrading capability of the 20 kDa polypeptide, 0.5 mg/ml of C3 (prepared according to Tack et al., *Meth. Enzymol.* 80:64-101, 1984) was copolymerized in a sodium dodecyl sulfate (SDS) gel-containing 15% acrylamide (15% SDS-PAGE gel). Pneumococcal supernatants were obtained from cultures of *S. pneumoniae* strain CP1200 grown to exponential phase in Todd Hewitt broth; pneumococcal lysates were obtained by incubating 5×10^8 cells with 5% SDS for 30 minutes at room temperature. The lysate was concentrated 10 fold using a Centricon filtration device with a 10,000 mw cutoff (Amicon, Beverly, MA). The samples were not heated before electrophoresis.

Samples of supernatants and lysates were added to the 15% C3-containing SDS-PAGE gels and electrophoresis was carried out at 4°C at 150 V until the dye front ran out. The gel was washed successively with 50 ml of 2.5% Triton X-100 in water (2 times, 10 minutes), 2.5 % Triton X-100 in 50 mM Tris-HCl, pH 7.4 (2 times, 10 minutes), and 50 mM Tris-HCl, pH 7.4 (2 times, 10 minutes) to remove SDS. After washes, 50 ml of 50 mM Tris-HCl, pH 7.4, was poured into dishes containing the gels, and the dishes were covered and incubated at 37°C for 1.5 hour and overnight (about 16 hours). The gels were stained with Coomassie blue for 10 minutes and destained totally.

Two lytic bands were visualized, one of which was about 20 kDa in size, against the dark blue background in both lysates and supernatant. C3 degrading activities in the pneumococcal lysates were observed after a 1.5 hour incubation at 37°C, while C3 degrading activities in the Pn supernatant were observed after an overnight incubation. Therefore, C3 degrading activities appeared to be mainly cell associated.

Example 3

The gene encoding the 20 kD polypeptide is conserved in a number of *S. pneumoniae* strains.

DNA was obtained from a variety of *S. pneumoniae* strains (Clinical isolates of Type 1, Type 3, LOO2 and LOO3 (type 3), Type 4, Type 14 and Laboratory isolates CP1200, WU2, R6X, 6303,109,110, JY1119, JY182, and JY53) and SEQ ID NO:3 was used as a probe to detect the presence of nucleic acid encoding the 20 kD polypeptide in DNA from these strains. Isolated chromosomal DNA was digested with *EcoRI* and separated by electrophoresis. The DNA was transferred to a solid support and hybridized to end-labeled SEQ ID NO:3 under the hybridization and washing conditions of 6X SSC, 5X Denhart's, 0.5% SDS, 100 µg/ml denatured fragmented salmon sperm DNA hybridized at 65 °C overnight and washed in 2X SSC, 1 time at room temperature for 10 minutes and in 2X SSC, 0.1% SDS 1 time at 65°C for 15 minutes followed by two washes in 0.2X SSC, 0.1% SDS for 3 minutes each at room temperature.

Results indicated that SEQ ID NO:3 hybridized identically in each of the DNA samples tested indicating that the polypeptide appears to be conserved among strains. In some strains, the DNA encoding the 20 kDa C3-degrading polypeptide appears to be part of a larger open reading frame of 2478 bp that
5 putatively encodes a 92 kDa polypeptide.

Example 4

Southern blot of *S. pneumoniae* DNA probe

Five ug samples of genomic DNA were obtained from 11 strains of *S. pneumoniae*. Each sample was digested with the restriction enzyme Kpn1. The
10 samples were subsequently loaded onto an agarose gel and resolved by electrophoresis. The samples contained in the gel were subsequently transferred to a Hybond-N+ membrane available from Amersham (Upsalla, Sweden) by capillary transfer. A 540 bp SmaI fragment from the open reading frame was
15 random primer labeled with P³² using a T⁷QuickPrime kit (Pharmacia, Piscataway, NJ) and purified from non-incorporated nucleotides using NucTrap column (Stragene, La Jolla, CA) and hybridized.

The hybridization conditions were 6XSSC, 5X Denhardt, 0.5% SDS, and 100 µg/ml fragmented and denatured salmon sperm DNA hybridized overnight
20 at 65°C and washed in 2X SSC, 0.1% SDS once at room temperature for about 10 minutes followed by 1 time at, 65°C for about 15 minutes followed by at least one wash in 0.2XSSC, 0.1% SDS at room temperature for at least 3-5 minutes. The blot demonstrated that the 20 kDa gene was present in all tested strains of *S. pneumoniae*.

25

Example 5

Two DNA primers were prepared from SEQ ID NO:1 and utilized to amplify the nucleotide sequence encoding the 20 kDa polypeptide from *S. pneumoniae* (serotype 3) genomic DNA. The first primer, a 5'-primer, SEQ ID
30 NO:6, includes an ATG start codon of the 5' end of the nucleotide sequence, inserts a NcoI site, and had an Ala residue inserted after the ATG start codon to maintain a correct reading frame. The second primer, a 3'-primer, SEQ ID

NO:7, includes a termination codon at the 3' end of the nucleotide sequence and inserts a BamH1 site.

5'-GGGGG CCA TGG CC TCA AGC CTT TTA CGT GAA TTG-3';
(SEQ ID NO:6)

5'-GGGGG GGA TCC CTA GCT ATA TGA GAT AAA CTT TCC
TGC T-3'; (SEQ ID NO:7)

10 The two primers were synthesized on an Applied Biosystems 380A DNA synthesizer (Foster City, CA) using reagents purchased from Glen Research (Sterling, VA). Amplifications were performed utilizing a Perkin Elmer Thermocycler (ABI) according to the manufacturer's directions. The identified PCR product was ligated into the TA tailed PCR cloning vector PCR2.1,
15 available from Invitrogen, Carlsbad, CA, and used to transform OneShot Top10F' competent cells (Invitrogen). Kanamycin resistant transformants were screened by restriction enzyme analysis of plasmid DNA prepared by alkaline lysis. An approximately 500 bp insert fragment was identified and subsequently excised with restriction enzymes Nco1 and BamH1. The 500 bp fragment was
20 purified from a low melting agarose gel, and subsequently ligated into the Nco1-BamH1 sites of the T7 promoted expression vector pET 28a, available from Novagen (Madison, WI).

The ligation mixture was subsequently transformed into Top10F'cells (Invitrogen), and the kanamycin resistant transformants were screened by
25 restriction enzyme analysis of plasmid DNA prepared by alkaline lysis. A recombinant plasmid (pLP505) was subsequently transformed into BL21 (Novagen) cells and grown in SOB media supplemented with 30 µg/ml kanamycin. Cells were grown to an O.D.₆₀₀ of 0.6, and were subsequently induced with 0.4mM IPTG (Boehringer Mannheim, Indianapolis, Indiana) for 2-
30 4 hours. Whole cell lysates were prepared and electrophoresed on a 14% SDS-PAGE gel. The gel was stained with Coomassie and the expression product was detected. The coomassie stained gel revealed a band between the 28 kDa and the

18 kDa molecular weight markers, and was determined to be approximately 20 kDa.

The DNA sequence of the insert in the recombinant pLP505 plasmid was obtained using the ABI 370A DNA sequencer. The DNA sequence was aligned with the DNA sequence of SEQ ID NO:1, using the Pustell DNA matrix plot feature of MacVector (Oxford Molecular Group, Campbell, CA). Alignment of the DNA sequence obtained from the pLP505 plasmid, SEQ ID NO:1, and the *S. pneumoniae* (serotype 4) genome, revealed that the open reading frame (ORF) that codes for the 20 kDa polypeptide may be part of a larger ORF, i.e., a 2478 bp in the serotype 4 genome, that codes for a polypeptide with a predicted MW of approximately 92 kDa (SEQ ID NO: 4). DNA SEQ ID NO:4 encodes for a predicted amino acid sequence as shown in SEQ ID NO:5.

The *S. pneumoniae* (serotype 4) genome sequence was obtained from The Institute for Genomic Research at www.tigr.org and/or through NCBI at www.ncbi.nlm.nih.gov, using the ClustalW feature of MacVector, (Oxford Molecular Group, Campbell CA). A sequence comparison was made between the 20 kDa amino acid sequence (SEQ ID NO:2) and the predicted 92 kDa amino acid sequence (SEQ ID NO:5).

Based upon the available genomic DNA (serotype 4) sequence, two primers flanking the 2478 bp ORF were designed and subsequently synthesized using the ABI 380A DNA synthesizer (SEQ ID NOS:8 and 9). SEQ ID NO:8 was an *S. pneumoniae* 5'-primer having an inserted NcoI site and a "Glu" residue added after the ATG start codon to maintain a correct reading frame. SEQ ID NO:9, was an *S. pneumoniae* 3'-primer having an inserted HindIII site.

5'- CCC GGG CCA TGG CTA AAA TTA ATA AAA AAT ATC TAG
-3'; (SEQ ID NO:8)

5'-CCG GGC AAG CTT TTA CTT ACT CTC CT-3'-; (SEQ ID NO:9)

An approximately 2400 bp DNA fragment was then amplified from the 4 different *S. pneumoniae* serotypes (serotype 3, 5, 6B and 7) resulting in 4

fragments. Each of the 4 fragments were subsequently ligated into the PCR cloning vector PCR2.1 (Invitrogen), and used to transform OneShot Top 10F' cells (Invitrogen). Kanamycin resistant transformants were screened by restriction analysis of the plasmid DNA prepared by alkaline lysis. A

- 5 recombinant plasmid containing the serotype 7 PCR product was identified, e.g., pLP512. The DNA sequence was obtained from the serotype 7 clone using the ABI model 370A DNA sequencer. The DNA sequence was essentially identical to SEQ ID NO:4 and encoded a predicted amino acid sequence essentially identical to SEQ ID NO:5.

10

Example 6

Western blot detection of 92 kDa Polypeptide in Whole Pneumococci

- Recombinant approximately 20 kDa C3 degrading polypeptide was
- 15 purified from *Escherichia coli* strain BLR containing plasmid pDF122. Plasmid pDF122 contains the polynucleotide shown in SEQ ID NO:1 expressed under control of the T7 phage promoter system. The bacterial cells were grown to mid-log phase in Hy-Soy/yeast Extract medium containing ampicillin to select for the plasmid. Expression of the recombinant polypeptide was induced by adding
- 20 IPTG to a concentration of 1mM and continuing incubation for an additional 3 hours. The bacterial cells were harvested by centrifugation and resuspended in Tris buffered saline, pH 7.2. Cells were mechanically lysed in a French Pressure cell and the insoluble material including inclusion bodies were pelleted in a centrifuge. The pellet was solubilized in 3 M Urea buffered with 100mM NaPO₄,
- 25 pH 8.0 containing 0.1% Triton X-100. After pelleting and discarding insoluble material (centrifuged at ~100,000 x g), the soluble r 20kDa polypeptide was exchanged into 0.1% Zwittergent 3-12 (Calbiochem-Behring) replacing the Urea and then into 100 mM NaPO₄ pH 8.0 replacing the detergent. SDS-PAGE analysis confirmed that the ~20 kDa material remained soluble. The His-tagged
- 30 recombinant polypeptide was dialyzed into 50 mM NaPO₄ buffer, pH 8.0 and absorbed onto a Ni column equilibrated with the same buffer. The column was sequentially washed with 50 mM NaPO₄ at pH 7.0, 6.6, and 5.5 until baseline absorbances (OD₂₈₀) were reached with each buffer. The bound ~r20 kDa

polypeptide was eluted with 50 mM NaPO₄, pH 4.5. The eluted polypeptide was approximately 90% homogeneous in SDS-PAGE analysis.

This polypeptide was used to immunize Swiss-Webster mice. Five µg doses of the polypeptide were mixed with 50 µg of Monophosphoryl lipid A (Ribi Immunochemicals) and injected intramuscularly into mice. Animals were immunized at weeks 0, 4, and 6 and exsanguinated at week 8. The sera were pooled together and found to contain high titered antisera against the immunogen.

Pneumococcal strains CP1200, T3, and T7 were grown in Todd-Hewitt broth containing yeast Extract and the cells pelleted by centrifugation. Pneumococcal cells were lysed in SDS-PAGE cracking buffer under reducing conditions by boiling for 5 minutes. The lysates were loaded onto a 10% SDS-PAGE gel and electrophoresed. Separated polypeptides were electroblotted onto Nitrocellulose and the filter blocked with 5% BLOTTO in phosphate buffered saline, pH 7.4. The polyclonal antisera was diluted 1:2000 in BLOTTO and used to probe the separated polypeptides. Bound antibodies were detected with Alkaline phosphatase conjugated goat anti-mouse IgG. The Western Blot is shown in Figure 8 and described further in the brief description of the figures.

20

Example 7

Evidence of that the 20 kDa and 92 kDa Polypeptides Degrade C3

The gel shown in Figure 9 is a Western blot. This data were obtained by incubating a 20 kDa polypeptide and a 92 kDa polypeptide with biotinylated and methylamine-treated C3 as shown in the table below. These polypeptides were expressed from a T7 vector.

METHYLAMINE-TREATED C3: 500 µl of purified human C3 at concentration of 4.46 mg/ml (prepared according to the method of Tack et al., Meth. Enzymol. 80:64-101, 1984) was incubated with 55 µl of 1 M methylamine (CH₃NH₂) in 0.1 M TRIS/0.01 M EDTA, pH 8.0 for 80 minutes at 37°C, then dialyzed overnight in 0.1 M TRIS/0.01 M EDTA, pH 8.0, at 4°C. The following day, the C3 was incubated again with 60 µl of 1 M methylamine in 0.1 M

TRIS/0.01 M EDTA for 80 minutes at 37°C, then dialyzed overnight in 50 mM NaHCO₃ at 4°C.

BIOTINYLATION of METHYLAMINE-TREATED C3: 2 mg of methylamine-treated C3, prepared above, was incubated with 75 µl of NSC-biotin (Pierce) made as 1 mg NSC biotin in 1 ml H₂O for 30 minutes at room temperature. The biotinylated C3 was dialyzed overnight in 50 mM NaHCO₃, pH 8.0 at 4°C.

Reagents	Tube A – 20 kDa	Tube C – 92 kDa
Polypeptide	34 µl 20 kDa polypeptide 5.6 x 10 ⁻⁹ moles	100 µl 92 kDa polypeptide 5.6 x 10 ⁻⁹ moles
Biotinylated C3	3 µl C3 71 x 10 ⁻¹² moles	3 µl C3 71 x 10 ⁻¹² moles
Buffer	63 µl 0.01% BSA/PBS	63 µl 0.01% BSA/PBS

10

Samples from Tubes A and C were removed after 66 hours of incubation, reduced by boiling for 2 minutes in reducing buffer with β mercaptoethanol, electrophoresed on 7.5% SDS-PAGE, and transferred to nitrocellulose paper for Western blotting. Western blotting was carried out for one hour according to standard procedures (Tobin et al. *PNAS USA* 76:4350-4354, 1979). Membranes were incubated with horseradish-peroxidase-avidin (1:10,000 dilution) and developed with the Supersignal CL-HRP Substrate System (Amersham) to detect biotinylated C3 fragments.

In Figure 9 lane 1 is C3 alone (control), lanes 2 and 3 are the 20 and 92 kDa polypeptides, and lanes 4 and 5 are again C3 alone (controls). Along the right margin the position of the α-chain of C3, the β-chain of C3, and the C3 fragment present in the control samples are shown. The position of a C3 fragment in the 20 kDa digest and several fragments in the 92 kDa digest are identified.

The second experiment used the reagents of the first (expressed polypeptides, biotinylated C3, and buffer) in the same proportions as shown in

the table above. C3 control samples were removed at time 0, 2 hours, 8 hours, and 24 hours. However, instead of doing a Western blot, the samples were reduced on 7.5% SDS-PAGE, reasoning that some additional degradation fragments might not be biotinylated and would therefore show up on a

5 Coomassie stained SDS-PAGE gel. Indeed, that is what was observed. Only the 20 (lane A8) and 92 (lane C8) kDa polypeptides were analyzed on this gel. The gel of Figure 10 shows along the right margin the position of the α -chain of C3 and the β -chain of C3, and the 92 kDa polypeptide in lane C8. In lane C8 are at least two fragments ranging in size from ~90 kDa to 75 kDa. There are also
10 several fragments beneath the β -chain at 75 kDa. In lane A8 is a major degradation fragment just below the β -chain of C3. Thus, both the 92 kDa polypeptide in lane C8 and the 20 kDa polypeptide in lane A8 degrade C3.

15 Example 8

Preparation of recombinant 92 kDa protein and generation of polyclonal antiserum

The insert in plasmid pLP512 (see Example 5) was excised with NcoI
20 and HindIII. A fragment of the expected size was purified from low melting point agarose, and subsequently ligated into the NcoI- HindIII sites of the T7 promoted expression vector pET28a (Novagen, Madison, WI).

The ligation mixture was subsequently transformed into Top10F' cells (Invitrogen) and the kanamycin resistant transformants were screened as
25 described previously in Example 5. A recombinant plasmid (pLP515) was subsequently transformed into BL21 cells (Novagen) and grown in SOB media supplemented with 30 μ g/ml kanamycin. Cells were grown to an O.D. ₆₀₀ of 0.6, and were subsequently induced with 0.4 mM IPTG (Boehringer Mannheim, Indianapolis, IN) for 2-4 hours. Whole cell lysates were prepared and
30 electrophoresed on a 10% SDS-PAGE gel. Coomassie staining of the gel revealed a new band of approximately 80 kDa as compared to a pre-induction sample.

Recombinant protein coded by the ~92 kDa ORF was purified from *E. coli* strain BL21 (Novagen) containing the plasmid pLP515. Bacterial cells were grown to mid log phase in SOB medium containing 50 µg/ml kanamycin to select for the plasmid. Expression of the recombinant polypeptide was induced by addition
5 IPTG to 0.4 mM and continuing incubation for 3 hours. The bacterial cells were harvested by centrifugation and resuspended in Tris-Buffered Saline, pH7.2. Cells were mechanically lysed in a French Pressure Cell and the insoluble material including inclusion bodies were pelleted in a centrifuge at 7700 x g at 4 degrees for 10 minutes. The pellet containing the inclusion bodies was
10 resolubilized and the soluble protein was purified by ion exchange chromatography. Recombinant protein was used to generate polyclonal antibodies in mice. Briefly, 5 µg of protein was adjuvanted for each dose with 20 µg QS21 and injected subcutaneously into the necks of a 6-8 weeks Swiss Webster mice. The mice were bled and vaccinated at week 0, vaccinated at week
15 4, then bled and exsanguinated at week 6. 10 mice were vaccinated with the recombinant 92 kDa protein adjuvanted with QS21. Pooled sera were used at a 1:1000 dilution to examine whole cell lysates and concentrated culture supernatants from several serotypes of *S. pneumoniae* on a Western Blot. The sera reacted specifically to a protein in whole cell lysates and concentrated
20 culture supernatants whose molecular weight was approximately 90 kDa.

Example 9

The chart of Figure 11 summarizes the results from an intranasal (IN)
25 challenge of CBA/CAHN *xid/J* mice vaccinated with r92 kDa protein, prepared as described in Example 8 (SEQ ID NO:5) adjuvanted with monophosphoryl lipid A (MPL).

10 mice per group were vaccinated subcutaneously in the neck region at weeks 0, 3, and 5, with either 5 µg each of 92 kDa adjuvanted with 50 µg MPL,
30 1 µg each of Type 3 capsule conjugated to the protein carrier CRM197 adjuvanted with 100 µg aluminum phosphate, or phosphate buffered saline (PBS) alone, in a sample volume of 10 µl. CRM 197 is a genetically detoxified version of diphtheria toxin. Each mouse was challenged in at week 7 with 1×10^6

cfu's of Type 3 *S. pneumoniae* in a 10 µl sample volume. Nasal tissue was isolated 3 days after challenge and the number of type 3 *S. pneumoniae* colony-forming units per gram of nasal tissue determined by plating on selective media. Type 3 is capsule isolated from *S. pneumoniae* serotype 3, and then conjugated to protein carrier CRM 197, through reductive animation (see U.S. Patent No. 5, 360,897 for preparation of the Type 3 control; see U.S. Patent No. 5,614,382 for genetically detoxified version of dithera toxin; see U.S. Patent 4,902,506 with regard to using CRM 197 as a carrier).

The chart in Figure 11 shows that in this model, both the negative controls, mice adjuvanted with MPL and naïve mice at 6 weeks of age, had a survival rate of approximately 30% when challenged intranasally, while the positive control, the Type 3 conjugate adjuvanted with aluminum phosphate, offered 100% protection against death to the end of the study, approximately 14 days. When mice immunized with r92 kDa were challenged, 100% survived to the end of the study, indicating that r92 kDa protein does offer protection against death from intranasal challenge by Type 3 *S. pneumoniae*.

Example 10

METHODS

Recombinant 92 kDa (r92 kDa) polypeptide (SEQ ID NO:5) were incubated with purified human C3 for 2 hours, 6 hours, and 26 hours at 37°C in the following ratios:

Tube A (Control): C3 - 3µl [7.2x10-12 moles] PBS/0.01% bovine serum albumin (BSA) - 100 µl; Tube B: r92 kDa - 50 µl [5.6x10-10 moles] C3 - 3 µl [7.2x10-12 moles] PBS/0.01% BSA - 50 µl.

At each time point (2, 6, and 26 hours), 20 µl samples were removed from Tubes A and B reducing buffer was added, and the samples were boiled at 100°C for 2 minutes.

After boiling, samples were electrophoresed on 7.5% SDS-PAGE under reducing conditions. The SDS-PAGE gel was stained with Coomassie blue. The resultant gel is shown in Figure 12.

Lane 1 - molecular weight standards (reading from top of gel: 200 kDa, 116 kDa, 97 kDa, 66 kDa, 45 kDa); Lanes 2 and 4 - 2 hour incubation; Lane 2 - C3 control (Tube A); Lane 4 - r92 kDa sample (Tube B); Lanes 5 and 7 - 6 hour incubation; Lane 5 - C3 control (Tube A); Lane 7 - r92 kDa sample (Tube B); Lanes 8 and 10 - 26 hour incubation; Lane 8 - C3 control (Tube A); Lane 10 - r92 kDa sample (Tube B).

INTERPRETATION of GEL

10 The α -chain of C3 runs at about 115 kDa. The β -chain of C3 runs at about 75 kDa. These are marked on the gel. The band at about 66 kDa is albumin. Lanes 2 and 4 are the 2-hour incubation. Compared to Lane 2 (C3 control), Lane 4 shows cleavage of the C3 α -chain - new fragment at ~97 kDa. Lanes 5 and 7 are the 6-hour incubation. There is the same new 97 kDa cleavage
15 fragment in Lane 7. Lanes 8 and 10 are the 26-hour incubation. Compared to Lane 8 (C3 control), Lane 10 shows continued cleavage of the α -chain.

 It will be appreciated by those skilled in the art that while the invention has been described above in connection with particular embodiments and
20 examples, the invention is not necessarily so limited and that numerous other embodiments, examples, uses, modifications and departures from the embodiments, examples and uses may be made without departing from the inventive scope of this application.

What is claimed is:

1. An isolated polypeptide comprising an amino acid sequence; wherein the nucleic acid sequence encoding the amino acid sequence of the isolated polypeptide hybridizes under highly stringent hybridization conditions to at least a portion of either (a) the nucleotides in SEQ ID NO:1 or a complementary strand thereof, or (b) the nucleotides in SEQ ID NO:4 or a complementary strand thereof.
2. The polypeptide of claim 1 which is isolated from *S. pneumoniae*.
3. The polypeptide of claim 1 which is a recombinant polypeptide.
4. The polypeptide of claim 1 having a molecular weight of about 15 kDa to about 25 kDa.
5. The polypeptide of claim 1 having a molecular weight of about 75 kDa to about 95 kDa.
6. The polypeptide of claim 1 which degrades human complement protein C3.
7. The polypeptide of claim 1 comprising at least 15 sequential amino acids.
8. An isolated polypeptide comprising at least 15 sequential amino acids of SEQ ID NO:2 or SEQ ID NO:5.
9. An isolated polypeptide comprising SEQ ID NO:2 or SEQ ID NO:5.

10. An immune system stimulating composition comprising an effective amount of the polypeptide of claim 1 and a therapeutically acceptable carrier.
11. The immune system stimulating composition of claim 10 wherein the polypeptide is isolated from *S. pneumoniae*.
12. The immune system stimulating composition of claim 11 further comprising at least one other immune system stimulating polypeptide isolated from *S. pneumoniae*.
13. The immune system stimulating composition of claim 10 wherein the polypeptide is effective to immunize or treat a mammalian subject against *S. pneumoniae* infection or colonization.
14. The immune system stimulating composition of claim 13 wherein the polypeptide is provided in an amount effective to provide a therapeutic effect to the mammalian subject.
15. An antibody capable of binding to the polypeptide of claim 1.
16. The antibody of claim 15 which is a monoclonal antibody.
17. The antibody of claim 16 which is obtained from a mouse, a rat, a goat, a chicken, a human, or a rabbit.
18. An isolated nucleic acid molecule which hybridizes under highly stringent hybridization conditions to at least a portion of either (a) the nucleotides in SEQ ID NO:1 or a complementary strand thereof, or (b) the nucleotides in SEQ ID NO:4 or a complementary strand thereof.
19. The nucleic acid molecule of claim 18 isolated from *S. pneumoniae*.

20. The nucleic acid molecule of claim 19 which encodes a polypeptide that degrades human complement C3.
21. A vector comprising the nucleic acid molecule of claim 18.
22. A cell comprising the nucleic acid of claim 18.
23. The cell of claim 22 which is a bacterium or a eukaryotic cell.
24. An isolated nucleic acid molecule comprising the nucleic acid sequence of SEQ ID NO:1 or its complementary strand or SEQ ID NO:4 or its complementary strand.
25. An RNA molecule transcribed by a double-stranded DNA sequence comprising SEQ ID NO:1 or SEQ ID NO:4.
26. A method for producing an immune response to *S. pneumoniae* in a mammal, the method comprising administering to a mammal a composition comprising:
 - (a) a therapeutically effective amount of the polypeptide of claim 1; and
 - (b) a pharmaceutically acceptable carrier.
27. The method of claim 26 wherein the immune response is a B cell response, a T cell response, an epithelial cell response, or an endothelial cell response.
28. The method of claim 26 wherein the polypeptide is at least 15 amino acids in length.

29. The method of claim 26 wherein the composition further comprises at least one other immune system stimulating polypeptide from *S. pneumoniae*.
30. A method for inhibiting *S. pneumoniae*-mediated C3 degradation, the method comprising contacting an *S. pneumoniae* bacterium with an antibody capable of binding to the polypeptide of claim 1.
31. A method for inhibiting C3-mediated inflammation and rejection in xenotransplantation, the method comprising expressing on the surface of an organ of an animal used in xenotransplantation the polypeptide of claim 1.
32. An isolated nucleic acid molecule having a nucleic acid sequence selected from the group consisting of SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, and SEQ ID NO:9.
33. An isolated polypeptide having at least 80% sequence identity to SEQ ID NO:2 or SEQ ID NO:5.
34. An isolated polypeptide of about 15 kDa to about 25 kDa from *S. pneumoniae* that degrades human complement protein C3.
35. An isolated polypeptide of about 75 kDa to about 95 kDa from *S. pneumoniae* that degrades human complement protein C3.

ATGTCAAGCCTTTTACGTGAATTGTATGCTAAACCCTTATCAGAACGCCATGTAGAATCTGATGGCC
TTATTTTCGACCCAGCGCAAATCACAAGTCGAACCGCCAATGGTGTTGCTGTACCGCACGGAGACC
ATTATCACTTTATTCTTATTACAACTGTACCTTTGGAAGAAAAATTGGCTCGTATTATTCCCCTTC
GTTATCGTTCAAACCAATTGGGTACCAGATTCAAGACCAGAACAACCAAGTCCACAATCGACTCCGG
AACCTAGTCCAAGTCCGCAACCTGCACCAAATCCTCAACCAGCTCCAAGCAATCCAATTGATGAGA
AATTGGTCAAAGAAGCTGTTGAAAAGTAGGCGATGGTTATGTCTTTGAGGAGAATGGAGTTCCTC
GTTATATCCCAGCCAAGGATCTTTCAGCAGAAACAGCAGCAGGCATTGATAGCAAACCTGGCCAAGC
AGGAAAGTTTATCTCATAAGCTGCAGTTAGATCCATTA

Figure 1 (SEQ ID NO: 1)

MSSLLRELYAKPLSERHVESDGLIFDPAQITSRTANGVAVPHGDHYHFIPYSQLSPLEEKLARIPLRYR
SNHWVPDSRPEQSPQSTPEPSPSPQPAPNPQPAPSNPIDEKLVKEAVRKVG DG YVFEENGVPRIIPA
KDLSAETAAGIDSKLAKQESLSHKLQLDPL

Figure 2 (SEQ ID NO: 2)

10 20 30 40 50 60 70
ATGTCAGGCCTTTTACGTGATTGTATGCTAAACCTTATCAGAACGCCATGTAGAACTCTGATGGCCTTATTTTC
TACAGTTTCGGAAATGCACTTAACATACGATTGGGAATAGTCTTGCGGTACATCTTAGACTACCGGAATAAAO
H S S L L R E L Y A K P L S E R H U E S D G L I F>

80 90 100 110 120 130 140 150
GACCCAGCGCAATCACAAGTCGAACCGCCAAATGGTGTGCTGTACCGCACGGAGACCATTATCACTTTATTCCT
CTGGGTGCGCTTATGTTGAGCTTGGCGTTACCAACACGCATGGCGTGCCTCTGGTAATAGTGAATAAGGA
D P A Q I T S A T A N G U A U P H G D H V H F I P>

160 170 180 190 200 210 220
TATTCACAACCTGTACCTTTGGAGGAAATTTGGCTCGTATTATTCCTTCCTTATCGTTCAAAACCATTTGGTA
ATAAGTGTGACAGTGGAAACCTTCTTTTAAACCGAGCATATAAGGGGAGCAATAGCAAGTTTGGTAACCCAT
Y S Q L S P L E E K L A R I I P L R Y R S N H W U>

230 240 250 260 270 280 290 300
CCAGATTCAAGACCAGAACACCAAGTCCACAATCGACTCCGGACCTAGTCCAACTCCGCACCTGCACCAAT
GGTCTAAGTTCTGGTCTTGTGTTGAGGTGTTAGCTGAGGCTTGGATCAGGTTTCAAGGCTTGGACGTGGTTA
P O S R P E Q P S P Q S T P E P S P S P Q P A P N>

310 320 330 340 350 360 370
CCTCAACCAAGCTCCAAAGCAATCCAAATGATGGAATTTGGTCAAGAGAGCTGTTCAAAAGTAGGCGATGTTAT
GGAATTTGGTCGAGGTTTGGTTAGTTTAACTACTCTTTAAACAGTTTCTTCGACAAAGCTTTTCATCCGCTACCAATA
P Q P A P S N P I D E K L U K E A U R K U G D G Y>

380 390 400 410 420 430 440 450
GTCTTTGAGGAGAAATGGAGTTCCTCGTTATATCCCAGCCAGGATCTTTCAGCAGAACAGCAGCAGGCATTGAT
CAGAACTCCTCTTACCTCAGGGAGCAATATAGGGTCGGTTCCTAGAAAGTCGTCTTTGTCGTGTCGGTAACTA
V F E E N G U P R Y I P A K D L S A E T A A G I D>

460 470 480 490 500
AGCAAACTGGCCAAAGCAGGAAAGTTTATCTCATAGCTGCAGTTAGATCCATTA
TCGTTTGACCGGTTCTGCTCTTCAATAGAGTATTCAGCTCAATCTAGGTAT
S K L A K Q E S L S H K L Q L D P L>


Figure 3
(SEQ ID NOS: 1, 2, and 3)

Fig. 4 (SEQ ID NO:4)

ATGAAAATTAATAAAAAATATCTAGCAGGTTTCAGTGGCAGTCCTTGCCCTAAGTGTTTGTTCCCTATGA
ACTTGGTTCGTACCAAGCTGGTCAGGTTAAGAAAGAGTCTAATCGAGTTTCTTATATAGATGGTGAT
CAGGCTGGTCAAAAGGCAGAAAACTTGACACCAGATGAAGTCAGTAAGAGGGAGGGGATCAACGC
CGAACAAATCGTCATCAAGATTACGGATCAAGGTTATGTGACCTCTCATGGAGACCATTATCATTAC
TATAATGGCAAGGTCCTTATGATGCCATCATCAGTGAAGAGCTCCTCATGAAAGATCCGAATTATC
AGTTGAAGGATTGAGACATTGTCAATGAAATCAAGGGTGGTTATGTTATCAAGGTAGATGGAAAATA
CTATGTTTACCTTAAGGATGCAGCTCATGCGGATAATATTGCGACAAAAGAAGAGATTAAACGTCAG
AAGCAGGAACACAGTCATAATCACGGGGGTGGTTCTAACGATCAAGCAGTAGTTGCAGCCAGAGC
CCAAGGACGCTATACAACGGATGATGGTTATATCTTCAATGCATCTGATATCATTGAGGACACGGGT
GATGCTTATATCGTTCCTCACGGCGACCATTAACATTACATTCCTAAGAATGAGTTATCAGCTAGCG
AGTTAGCTGCTGCAGAAGCCTATTGGAATGGGAAGCAGGGATCTCGTCCTTCTTCAAGTTCTAGTTA
TAATGCAAATCCAGCTCAACCAAGATTGTCAGAGAACCACAATCTGACTGTCACTCCAACCTTATCAT
CAAAATCAAGGGGAAAACATTTCAAGCCTTTTACGTGAATTGTATGCTAAACCCTTATCAGAACGCC
ATGTGGAATCTGATGGCCTTATTTTCGACCCAGCGCAAATCACAAGTCGAACCGCCAGAGGTGTAG
CTGTCCCTCATGGTAACCATTACCACTTTATCCCTTATGAACAAATGTCTGAATTGGAAAAACGAATT
GCTCGTATTATCCCCTTCGTTATCGTTCAAACCATTGGGTACCAGATTCAAGACCAGAACAACCAA
GTCCACAATCGACTCCGGAACCTAGTCCAAGTCCGCAACCTGCACCAAATCCTCAACCAGCTCCAA
GCAATCCAATTGATGAGAAATTGGTCAAAGAAGCTGTTGAAAAGTAGGCGATGGTTATGTCTTTGA
GGAGAATGGAGTTTCTCGTTATATCCCAGCCAAGGATCTTTCAGCAGAAACAGCAGCAGGCATTGA
TAGCAAACCTGGCCAAGCAGGAAAGTTTATCTCATAAGCTAGGAGCTAAGAAAACCTGACCTCCCATCT
AGTGATCGAGAATTTTACAATAAGGCTTATGACTTACTAGCAAGAATTCACCAAGATTTACTTGATAA
TAAAGGTCGACAAGTTGATTTTGAGGCTTTGGATAACCTGTTGGAACGACTCAAGGATGTCCCAAGT
GATAAAGTCAAGTTAGTGGATGATATTCTTGCCCTTCTAGCTCCGATTCTGCATCCAGAACGTTTAG
GAAAACCAAATGCGCAAATTACCTACACTGATGATGAGATTCAAGTAGCCAAGTTGGCAGGCAAGT
ACACAACAGAAGACGGTTATATCTTTGATCCTCGTGATATAACCAAGTATGAGGGGGATGCCTATGT
AACTCCACATATGACCCATAGCCACTGGATTAATAAAGATAGTTTGTCTGAAGCTGAGAGAGCGGC
AGCCCAGGCTTATGCTAAAGAGAAAAGGTTTGACCCCTCCTTCGACAGACCATCAGGATTCAGGAAA
TACTGAGGCAAAAAGGAGCAGAAGCTATCTACAACCGCGTGAAAGCAGCTAAGAAGGTGCCACTTGA
TCGTATGCCTTACAATCTTCAATATACTGTAGAAGTCAAAAACGGTAGTTTAATCATACCTCATTATG
ACCATTACCATAACATCAAAATTTGAGTGGTTTGACGAAGGCCTTTATGAGGCACCTAAGGGGTATAC
TCTTGAGGATCTTTTGGCGACTGTCAAGTACTATGTGCAACATCCAAACGAACGTCCGCATTGAT
AATGGTTTTGGTAACGCTAGCGACCATGTTCAAAGAAACAAAAATGGTCAAGCTGATACCAATCAA
CGGAAAAACCAAGCGAGGAGAAACCTCAGACAGAAAAACCTGAGGAAGAAACCCCTCGAGAAGAG
AAACCGCAAAGCGAGAAACAGAGTCTCAAACCAACAGAGGAACCAGAAGAATCACCAGAGGA
ATCAGAAGAACCTCAGGTCGAGACTGAAAAGGTTGAAGAAAAACTGAGAGAGGCTGAAGATTTACT
TGAAAAAATCCAGGATCCAATTATCAAGTCCAATGCCAAAGAGACTCTCACAGGATTAAAAAATAATT
TACTATTTGGCACCCAGGACAACAATACTATTATGGCAGAAGCTGAAAAACTATTGGCTTTATTAAG
GAGAGTAAG

Fig. 5 (SEQ ID NO: 5)

MKINKKYLAGSVAVLALSVCSYELGRHQAGQVKKESNRVSYIDGDQAGQKAENLTPDEVSKREGINA
EQIVIKITDQGYVTSHGDHYHYNGKVPYDAIISEELLMKDPNYQLKDS DIVNEIKGGYVIKVDGKYV
YLKDAAHADNIRTKEEIKRQKQEHSHNHGGGSNDQAVVAARAQGRYTTDDGYIFNASDIIEDTGDAYI
VPHGDHYHYIPKNELSASELAAAEAYWNGKQGSRPSSSSSYNANPAQPRLSENHNLTVTPTYHQNQ
ENISSLLRELYAKPLSERHVESDGLIFDPAQITSRTARGVAVPHGNHYHFIPYEQMSELEKRIARIPLRY
RSNHWVPDSRPEQSPQSTPEPSPSPQPAPNPQPAPSNPIDELVKEAVRKVGDGYVFEENGVSRYIP
AKDLSAETAAGIDSKLAKQESLSHKLGAKKTDLPSSDREFYNKAYDLLARIHQDLLDNKGRQVDFEAL
DNLLERLKDVP SDKVKLVDDILAFLAPIRHPERLGKPN AQITYTDD EIQVAKLAGKYTTEDGYIFDPRDI
TSDEGDAYVTPHMTSHWIKKDSLSEAERAAAQAYAKEKGLTPPSTDHQDSGNTEAKGAEAIYNRVK
AAKKVPLDRMPYNLQYTVEVKNGSLIIPHYDHYHNIKFEWFDEGLYEAPKGYTTLEDLLATVKYYVEHP
NERPHSDNGFGNASDHVQRNKNQADTNQTEKPSEEKPQTEKPEEETPREEKPQSEKPESPKPTEE
PEESPEESEEPQVETEKVEEKLREAEDLLGKIQDPIIKSNAKETLTGLKNNLLFGTQDNNTIMAEAEKL
LALLKESK



(SEQ ID NOS: 1 and 4)

Fig 6




Fig. 6 contd.
(SEQ ID NOS: 1 and 4)




Fig. 7
(SEQ ID NO:2 and a portion of
SEQ ID NO:5)




Figure 8




Fig. 9





Fig. 10



IN = Intranasa1

Fig. 11




Fig. 12

SEQUENCE LISTING

<110> HOSTETTER, Margaret K.
 FINKEL, David J.
 CHENG, Qi
 MASI, Amy W.
 REGENTS OF THE UNIVERSITY OF MINNESOTA
 AMERICAN CYNAMID COMPANY

<120> HUMAN COMPLEMENT C3-DEGRADING PROTEINASE FROM
 STREPTOCOCCUS PNEUMONIAE

<130> 11000570230

<140> Not Assigned

<141> 1999-09-24

<150> 60/101,736

<151> 1998-09-24

<150> 09/283,094

<151> 1999-03-31

<160> 10

<170> PatentIn Ver. 2.0

<210> 1

<211> 504

<212> DNA

<213> Streptococcus pneumoniae

<400> 1

atgtcaagcc	ttttacgtga	attgtatgct	aaacccttat	cagaacgcca	tgtagaatct	60
gatggcctta	ttttcgaccc	agcgcaaata	acaagtcgaa	ccgccaatgg	tgttgctgta	120
ccgcacggag	accattatca	ctttattcct	tattcacaac	tgtaaccttt	ggaagaaaaa	180
ttggctcgta	ttattcccct	tcgttatcgt	tcaaaccatt	gggtaccaga	ttcaagacca	240
gaacaaccaa	gtccacaatc	gactccggaa	cctagtccaa	gtccgcaacc	tgacacaaat	300
cctcaaccag	ctccaagcaa	tccaattgat	gagaaattgg	tcaaagaagc	tgttcgaaaa	360
gtaggcgatg	gttatgtctt	tgaggagaat	ggagttcctc	gttatatccc	agccaaggat	420
ctttcagcag	aaacagcagc	aggcattgat	agcaaactgg	ccaagcagga	aagtttatct	480
cataagctgc	agttagatcc	atta				504

<210> 2

<211> 168

<212> PRT

<213> Streptococcus pneumoniae

<400> 2

Met	Ser	Ser	Leu	Leu	Arg	Glu	Leu	Tyr	Ala	Lys	Pro	Leu	Ser	Glu	Arg
1				5					10					15	
His	Val	Glu	Ser	Asp	Gly	Leu	Ile	Phe	Asp	Pro	Ala	Gln	Ile	Thr	Ser
			20					25					30		
Arg	Thr	Ala	Asn	Gly	Val	Ala	Val	Pro	His	Gly	Asp	His	Tyr	His	Phe
		35					40					45			
Ile	Pro	Tyr	Ser	Gln	Leu	Ser	Pro	Leu	Glu	Glu	Lys	Leu	Ala	Arg	Ile
	50					55					60				

Ile Pro Leu Arg Tyr Arg Ser Asn His Trp Val Pro Asp Ser Arg Pro
 65 70 75 80

Glu Gln Pro Ser Pro Gln Ser Thr Pro Glu Pro Ser Pro Ser Pro Gln
 85 90 95

Pro Ala Pro Asn Pro Gln Pro Ala Pro Ser Asn Pro Ile Asp Glu Lys
 100 105 110

Leu Val Lys Glu Ala Val Arg Lys Val Gly Asp Gly Tyr Val Phe Glu
 115 120 125

Glu Asn Gly Val Pro Arg Tyr Ile Pro Ala Lys Asp Leu Ser Ala Glu
 130 135 140

Thr Ala Ala Gly Ile Asp Ser Lys Leu Ala Lys Gln Glu Ser Leu Ser
 145 150 155 160

His Lys Leu Gln Leu Asp Pro Leu
 165

<210> 3
 <211> 504
 <212> DNA
 <213> Streptococcus pneumoniae

<400> 3
 taatggatct aactgcagct tatgagataa actttcctgc ttggccagtt tgctatcaat 60
 gcctgctgct gtttctgctg aaagatcctt ggctgggata taacgaggaa ctccattctc 120
 ctcaaagaca taaccatcgc ctacttttctg aacagcttct ttgaccaatt tctcatcaat 180
 tggattgctt ggagctgggt gaggatttgg tgcaggttgc ggacttggac taggttccgg 240
 agtcgattgt ggacttgggt gttctgggtct tgaatctggg acccaatggg ttgaacgata 300
 acgaagggga ataatacgag ccaatttttc ttccaaaggt gacagttgtg aataaggaat 360
 aaagtgataa tgggtctccgt gcggtacagc aacaccattg gcggttcgac ttgtgatttg 420
 cgctgggtcg aaaataaggc catcagattc tacatggcgt tctgataagg gtttagcata 480
 caattcacgt aaaaggcttg acat 504

<210> 4
 <211> 2478
 <212> DNA
 <213> Streptococcus pneumoniae

<400> 4
 atgaaaatta ataaaaaata tctagcaggt tcagtgagcag tccttgccct aagtgtttgt 60
 tcctatgaac ttggctgctca ccaagctggg caggttaaga aagagtctaa tcgagtttct 120
 tatatagatg gtgatcaggc tgggtcaaaag gcagaaaact tgacaccaga tgaagtcagt 180
 aagagggagg ggatcaacgc cgaacaaatc gtcacaaaga ttacggatca aggttatgtg 240
 acctctcatg gagaccatta tcattactat aatggcaagg tcccttatga tgccatcatc 300
 agtgaagagc tcctcatgaa agatccgaat tatcagttga aggattcaga cattgtcaat 360
 gaaatcaagg gtggttatgt tatcaaggta gatggaaaat actatgttta ccttaaggat 420
 gcagctcatg cggataatat tcggacaaaa gaagagatta aacgtcagaa gcaggaacac 480
 agtcataatc acgggggtgg ttctaacgat caagcagtag ttgcagccag agcccaagga 540
 cgctatacaa cggatgatgg ttatatcttc aatgcattctg atatcattga ggacacgggt 600
 gatgcttata tcgttctctca cggcgaccat taccattaca ttccaaagaa tgagttatca 660
 gctagcgagt tagctgctgc agaagcctat tggaaatggga agcagggatc tcgtccttct 720
 tcaagttcta gttataatgc aaatccagct caaccaagat tgtcagagaa ccacaatctg 780
 actgtcactc caacttatca tcaaaatcaa ggggaaaaca tttcaagcct tttacgtgaa 840
 ttgtatgcta aacccttatc agaacgccat gtggaatctg atggccttat tttcgacca 900
 gcgcaaatca caagtcgaac cgccagaggt gtagctgtcc ctcatggtaa ccattaccac 960
 tttatccctt atgaacaaat gtctgaattg gaaaacagaa ttgctcgtat tattccctt 1020
 cgttatcggt caaaccattg ggtaccagat tcaagaccag aaccaaccaag tccacaatcg 1080
 actccggaac ctagtccaag tccgcaacat gcaccaaatc ctcaaccagc tccaagcaat 1140

```

ccaattgatg agaaattggt caaagaagct gttcgaaaag taggcgatgg ttatgtcttt 1200
gaggagaatg gagtttctcg ttatatccca gccaaaggatc tttcagcaga aacagcagca 1260
ggcattgata gcaaactggc caagcaggaa agtttatctc ataagctagg agctaagaaa 1320
actgacctcc catctagtga tcgagaattt tacaataagg cttatgactt actagcaaga 1380
attcaccaag atttacttga taataaagggt cgacaagttg attttgaggc tttggataac 1440
ctgttggaac gactcaagga tgtcccaagt gataaagtca agttagtgga tgatattctt 1500
gccttcttag ctccgattcg tcatccagaa cgtttaggaa aaccaaatgc gcaaattacc 1560
tacactgatg atgagattca agtagccaag ttggcaggca agtacacaac agaagacggg 1620
tatatctttg atcctcgtga tataaccagt gatgaggggg atgcctatgt aactccacat 1680
atgacccata gccactggat taaaaaagat agtttgtctg aagctgagag agcggcagcc 1740
caggcttatg ctaaagagaa aggtttgacc cctccttcga cagaccatca ggattcagga 1800
aatactgagg caaaaggagc agaagctatc tacaaccgag tgaaagcagc taagaagggtg 1860
ccacttgatc gtatgcctta caatcttcaa tatactgtag aagtcaaaaa cggtagttta 1920
atcatacctc attatgacca ttaccataac atcaaatttg agtggtttga cgaaggcctt 1980
tatgaggcac ctaaggggta tactcttgag gatctttttg cgactgtcaa gtactatgtc 2040
gaacatccaa acgaacgtcc gcattcagat aatgggtttg gtaacgctag cgaccatggt 2100
caaagaaaca aaaatggtca agctgatacc aatcaaacgg aaaaaccaag cgaggagaaa 2160
cctcagacag aaaaacctga ggaagaaacc cctcgagaag agaaaccgca aagcgagaaa 2220
ccagagtctc caaaaccaac agaggaacca gaagaatcac cagaggaatc agaagaacct 2280
caggctcgaga ctgaaaagggt tgaagaaaaa ctgagagagg ctgaagattt acttggaaaa 2340
atccaggatc caattatcaa gtccaatgcc aaagagactc tcacaggatt aaaaaataat 2400
ttactatttg gcacccagga caacaatact attatggcag aagctgaaaa actattgggt 2460
ttattaaagg agagtaag                                2478

```

<210> 5

<211> 826

<212> PRT

<213> Streptococcus pneumoniae

<400> 5

Met Lys Ile Asn Lys Lys Tyr Leu Ala Gly Ser Val Ala Val Leu Ala
 1 5 10 15

Leu Ser Val Cys Ser Tyr Glu Leu Gly Arg His Gln Ala Gly Gln Val
 20 25 30

Lys Lys Glu Ser Asn Arg Val Ser Tyr Ile Asp Gly Asp Gln Ala Gly
 35 40 45

Gln Lys Ala Glu Asn Leu Thr Pro Asp Glu Val Ser Lys Arg Glu Gly
 50 55 60

Ile Asn Ala Glu Gln Ile Val Ile Lys Ile Thr Asp Gln Gly Tyr Val
 65 70 75 80

Thr Ser His Gly Asp His Tyr His Tyr Tyr Asn Gly Lys Val Pro Tyr
 85 90 95

Asp Ala Ile Ile Ser Glu Glu Leu Leu Met Lys Asp Pro Asn Tyr Gln
 100 105 110

Leu Lys Asp Ser Asp Ile Val Asn Glu Ile Lys Gly Gly Tyr Val Ile
 115 120 125

Lys Val Asp Gly Lys Tyr Tyr Val Tyr Leu Lys Asp Ala Ala His Ala
 130 135 140

Asp Asn Ile Arg Thr Lys Glu Glu Ile Lys Arg Gln Lys Gln Glu His
 145 150 155 160

Ser His Asn His Gly Gly Gly Ser Asn Asp Gln Ala Val Val Ala Ala
 165 170 175

Arg Ala Gln Gly Arg Tyr Thr Thr Asp Asp Gly Tyr Ile Phe Asn Ala
 180 185 190
 Ser Asp Ile Ile Glu Asp Thr Gly Asp Ala Tyr Ile Val Pro His Gly
 195 200 205
 Asp His Tyr His Tyr Ile Pro Lys Asn Glu Leu Ser Ala Ser Glu Leu
 210 215 220
 Ala Ala Ala Glu Ala Tyr Trp Asn Gly Lys Gln Gly Ser Arg Pro Ser
 225 230 235 240
 Ser Ser Ser Ser Tyr Asn Ala Asn Pro Ala Gln Pro Arg Leu Ser Glu
 245 250 255
 Asn His Asn Leu Thr Val Thr Pro Thr Tyr His Gln Asn Gln Gly Glu
 260 265 270
 Asn Ile Ser Ser Leu Leu Arg Glu Leu Tyr Ala Lys Pro Leu Ser Glu
 275 280 285
 Arg His Val Glu Ser Asp Gly Leu Ile Phe Asp Pro Ala Gln Ile Thr
 290 295 300
 Ser Arg Thr Ala Arg Gly Val Ala Val Pro His Gly Asn His Tyr His
 305 310 315 320
 Phe Ile Pro Tyr Glu Gln Met Ser Glu Leu Glu Lys Arg Ile Ala Arg
 325 330 335
 Ile Ile Pro Leu Arg Tyr Arg Ser Asn His Trp Val Pro Asp Ser Arg
 340 345 350
 Pro Glu Gln Pro Ser Pro Gln Ser Thr Pro Glu Pro Ser Pro Ser Pro
 355 360 365
 Gln Pro Ala Pro Asn Pro Gln Pro Ala Pro Ser Asn Pro Ile Asp Glu
 370 375 380
 Lys Leu Val Lys Glu Ala Val Arg Lys Val Gly Asp Gly Tyr Val Phe
 385 390 395 400
 Glu Glu Asn Gly Val Ser Arg Tyr Ile Pro Ala Lys Asp Leu Ser Ala
 405 410 415
 Glu Thr Ala Ala Gly Ile Asp Ser Lys Leu Ala Lys Gln Glu Ser Leu
 420 425 430
 Ser His Lys Leu Gly Ala Lys Lys Thr Asp Leu Pro Ser Ser Asp Arg
 435 440 445
 Glu Phe Tyr Asn Lys Ala Tyr Asp Leu Leu Ala Arg Ile His Gln Asp
 450 455 460
 Leu Leu Asp Asn Lys Gly Arg Gln Val Asp Phe Glu Ala Leu Asp Asn
 465 470 475 480
 Leu Leu Glu Arg Leu Lys Asp Val Pro Ser Asp Lys Val Lys Leu Val
 485 490 495
 Asp Asp Ile Leu Ala Phe Leu Ala Pro Ile Arg His Pro Glu Arg Leu
 500 505 510

Gly Lys Pro Asn Ala Gln Ile Thr Tyr Thr Asp Asp Glu Ile Gln Val
 515 520 525
 Ala Lys Leu Ala Gly Lys Tyr Thr Thr Glu Asp Gly Tyr Ile Phe Asp
 530 535 540
 Pro Arg Asp Ile Thr Ser Asp Glu Gly Asp Ala Tyr Val Thr Pro His
 545 550 555 560
 Met Thr His Ser His Trp Ile Lys Lys Asp Ser Leu Ser Glu Ala Glu
 565 570 575
 Arg Ala Ala Ala Gln Ala Tyr Ala Lys Glu Lys Gly Leu Thr Pro Pro
 580 585 590
 Ser Thr Asp His Gln Asp Ser Gly Asn Thr Glu Ala Lys Gly Ala Glu
 595 600 605
 Ala Ile Tyr Asn Arg Val Lys Ala Ala Lys Lys Val Pro Leu Asp Arg
 610 615 620
 Met Pro Tyr Asn Leu Gln Tyr Thr Val Glu Val Lys Asn Gly Ser Leu
 625 630 635 640
 Ile Ile Pro His Tyr Asp His Tyr His Asn Ile Lys Phe Glu Trp Phe
 645 650 655
 Asp Glu Gly Leu Tyr Glu Ala Pro Lys Gly Tyr Thr Leu Glu Asp Leu
 660 665 670
 Leu Ala Thr Val Lys Tyr Tyr Val Glu His Pro Asn Glu Arg Pro His
 675 680 685
 Ser Asp Asn Gly Phe Gly Asn Ala Ser Asp His Val Gln Arg Asn Lys
 690 695 700
 Asn Gly Gln Ala Asp Thr Asn Gln Thr Glu Lys Pro Ser Glu Glu Lys
 705 710 715 720
 Pro Gln Thr Glu Lys Pro Glu Glu Glu Thr Pro Arg Glu Glu Lys Pro
 725 730 735
 Gln Ser Glu Lys Pro Glu Ser Pro Lys Pro Thr Glu Glu Pro Glu Glu
 740 745 750
 Ser Pro Glu Glu Ser Glu Glu Pro Gln Val Glu Thr Glu Lys Val Glu
 755 760 765
 Glu Lys Leu Arg Glu Ala Glu Asp Leu Leu Gly Lys Ile Gln Asp Pro
 770 775 780
 Ile Ile Lys Ser Asn Ala Lys Glu Thr Leu Thr Gly Leu Lys Asn Asn
 785 790 795 800
 Leu Leu Phe Gly Thr Gln Asp Asn Asn Thr Ile Met Ala Glu Ala Glu
 805 810 815
 Lys Leu Leu Ala Leu Leu Lys Glu Ser Lys
 820 825

<210> 6

<211> 34

<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<220>
<223> Incorporates a NcoI site and a DNA codon for ala.

<400> 6
ggggggccatg gcctcaagcc ttttacgtga attg 34

<210> 7
<211> 39
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<220>
<223> Incorporates a BamHI site.

<400> 7
ggggggggatc cctagctata tgagataaac tttcctgct 39

<210> 8
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<220>
<223> Incorporates a NcoI site and a DNA codon for Glu.

<400> 8
cccgggccat ggctaaaatt aataaaaaat atctag 36

<210> 9
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<220>
<223> Incorporates a HindD III site.

<400> 9
ccgggcaagc ttttacttac tctcct 26

<210> 10
<211> 40
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Probe

<220>

<223> Oligonucleotide

<400> 10.

gaaaacaata atgtagaaga ctactttaaa gaaggtaga

40

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/57 C12N9/52 C07K16/40 A01K67/027 A61K38/48
A61P37/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K A01K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 18930 A (HUMAN GENOME SCIENCES INC ;CHOI GIL H (US); HROMOCKYJ ALEX (US); J) 7 May 1998 (1998-05-07) SEQ ID NOs 55, 56, 65 & 66 page 98; claims 1-21 ---	1-31, 33-35
X	WO 98 18931 A (DOUGHERTY BRIAN A ;HUMAN GENOME SCIENCES INC (US); ROSEN CRAIG A () 7 May 1998 (1998-05-07) SEQ ID NO 94, 243, 258 claims 6-20 --- -/--	1-31, 33-35

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

3 February 2000

Date of mailing of the international search report

24.02.00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

van Klompenburg, W

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ANGEL ET AL: "Degradation of C3 by Streptococcus pneumoniae" JOURNAL OF INFECTIOUS DISEASES, US, CHICAGO, IL, vol. 170, no. 3, 1 January 1994 (1994-01-01), pages 600-608, XP002092872 ISSN: 0022-1899	34, 35
A	page 607, column 1; figures 3, 6-8 ---	1-33
X	NANDIWADA L S ET AL: "Genetic analysis of a C3 degrading proteinase in Streptococcus pneumoniae" ABSTRACTS OF THE GENERAL MEETING OF THE AMERICAN SOCIETY FOR MICROBIOLOGY, US, AMERICAN SOCIETY FOR MICROBIOLOGY, WASHINGTON, 1 January 1996 (1996-01-01), page 177 XP002076135 ISSN: 0067-2777	34
A	abstract ---	1-33, 35
A	WO 97 26008 A (UNIV MINNESOTA ; CLEARY PAUL P (US)) 24 July 1997 (1997-07-24) page 3, line 30 - page 5, line 2; claims 1-21; examples 1, 4, 6 ---	1-35
L, P, X	WO 99 15675 A (GREEN BRUCE A ; CHENG QI (US); FINKEL DAVID J (US); MASI AMY W (US)) 1 April 1999 (1999-04-01) claims 1-61; figures 1-7; examples 1-5 ---	1-35
P, X	WO 98 48022 A (NANDIWADA LAKSHMI S ; DUNNY GARY (US); UNIV MINNESOTA (US); HOSTETT) 29 October 1998 (1998-10-29) page 4, line 6 - line 10; claim 43; examples 1-5 -----	34

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 99/22362

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 26-30 are directed to a method of treatment of the human/animal body, and claim 31 at least partially, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on it family members

International Application No.

PCT/US 99/22502

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9818930	A	07-05-1998	AU 5194598 A	22-05-1998
			AU 6909098 A	22-05-1998
			EP 0942983 A	22-09-1999
			EP 0941335 A	15-09-1999
			WO 9818931 A	07-05-1998

WO 9818931	A	07-05-1998	AU 5194598 A	22-05-1998
			AU 6909098 A	22-05-1998
			EP 0942983 A	22-09-1999
			EP 0941335 A	15-09-1999
			WO 9818930 A	07-05-1998

WO 9726008	A	24-07-1997	US 5846547 A	08-12-1998
			AU 705732 B	27-05-1999
			AU 1582897 A	11-08-1997
			CA 2243755 A	24-07-1997
			EP 0877624 A	18-11-1998

WO 9915675	A	01-04-1999	AU 9510598 A	12-04-1999

WO 9848022	A	29-10-1998	AU 7156698 A	13-11-1998
